

MEDICAL REPORT

FOR THE YEAR

1930

GLASGOW ROYAL MATERNITY
AND WOMEN'S HOSPITAL



22501562733

Glasgow Royal Maternity and Women's Hospital

MEDICAL REPORT

For the Year 1930

Prepared by

H. R. MACLENNAN, M.B., Ch.B., Registrar to
the Hospital

GLASGOW :

AIRD & COGHILL, LTD., 24 Douglas Street, C.2

1931

WILLOW

PROPERTY

Ann Rep

WQ28

.BS2

G54

1930

CONTENTS.

	PAGE
Medical Staff, - - - - -	4
Foreword, - - - - -	5
Number and Classification of Patients, - - - - -	6
Numerical Summary of Cases Delivered in Hospital or Admitted after Delivery, - - - - -	9
Ante-natal Cases—Indoor, - - - - -	14
Obstetrical Cases—Indoor, - - - - -	15
Malpresentations—	
1. Vertex—Occipito-posterior, - - - - -	15
2. Breech, - - - - -	16
3. Face and Brow, - - - - -	18
4. Transverse, - - - - -	19
Multiple Pregnancy—	
1. Twins, - - - - -	20
2. Triplets, - - - - -	22
Obstetrical Operations—	
1. Forceps, - - - - -	23
2. Induction, - - - - -	24
3. Cæsarean Section, - - - - -	24
4. Version, - - - - -	29
5. Craniotomy, - - - - -	29
6. Oophorectomy, Salpingectomy, &c., - - - - -	31
7. Hysterectomy, - - - - -	31
8. Hysterotomy and Laparotomy, - - - - -	31
Contracted Pelvis, - - - - -	32
Ante-partum Hæmorrhage—	
1. Placenta Prævia, - - - - -	48, 52
2. Accidental Hæmorrhage, - - - - -	50, 57
Toxæmia of Pregnancy, - - - - -	64
Pyelitis, - - - - -	85
Heart Disease, - - - - -	89
Maternal Deaths, - - - - -	94
Stillbirths and Neonatal Deaths, - - - - -	107
Puerperal Pyrexia, - - - - -	107
Abortions, - - - - -	108
Mole, - - - - -	108
Appendix I.—	
1. Stobhill Cases, - - - - -	109
2. District Cases, - - - - -	109

MEDICAL STAFF.

Honorary Consulting Obstetric Physicians.

W. L. REID, M.D.

A. W. RUSSELL, M.B., C.M.

E. H. LAWRENCE OLIPHANT, M.D.

A. N. M'LELLAN, M.B., C.M.

R. JARDINE, M.D.

Honorary Consulting Physician—W. MACLENNAN, M.B., Ch.B.

Honorary Consulting Ophthalmic Surgeon—A. J. BALLANTYNE, M.D.

Obstetrical Units.

A.

Chief Obstetric Surgeon—Professor J. M. MUNRO KERR, M.D.

Visiting Obstetric Surgeon—D. M'INTYRE, M.D., F.R.C.S.E.

Assistant Obstetric Surgeons—A. M. HUNTER, M.B., Ch.B.

D. F. ANDERSON, M.B., Ch.B.

Extra Assistant Obstetric Surgeon—W. G. MACKAY, M.B., Ch.B.

B.

Chief Obstetric Surgeon—S. J. CAMERON, M.B., Ch.B.

Visiting Obstetric Surgeon—R. A. LENNIE, M.D.

Assistant Obstetric Surgeons—A. BARR, M.B., Ch.B.

E. D. MORTON, M.B., Ch.B.

Extra Assistant Obstetric Surgeon—W. C. ARMSTRONG, M.B., Ch.B.

C.

Chief Obstetric Surgeon—Professor J. HENDRY, M.A., B.Sc., M.B., Ch.B.

Visiting Obstetric Surgeon—J. HEWITT, M.B., Ch.B.

Assistant Obstetric Surgeons—D. BAIRD, M.B., Ch.B., B.Sc.

E. JACKSON, M.B., Ch.B., F.R.C.S.E.

Extra Assistant Obstetric Surgeon—A. FLEMING, M.D, B.Sc.

Visiting Anæsthetists—R. G. Grieve, M.B., Ch.B.

A. PILLANS, M.B., Ch.B.

Assistant Obstetric Surgeon (unattached)—A. MACARTNEY, M.B., Ch.B.

Senior Resident Obstetric Officer and Registrar—H. R. MACLENNAN, M.B., Ch.B.

Honorary Physician for Infant Ailments—S. GRAHAM, M.D.

Assistant to Honorary Physician for Infant Ailments—E. D. MORTON, M.B., Ch.B.

Acting Pathologist—D. BAIRD, M.B., Ch.B., B.Sc.

Medical Officer to V.D. Clinic—N. T. WATTIE, M.B., Ch.B., D.P.H.

Six Resident Obstetric Officers.

FOREWORD.

THE Fifth Annual Medical Report of the Glasgow Royal Maternity and Women's Hospital gives a brief record of the medical work carried out in the Hospital during the year commencing 1st January, 1930, and ending 31st December, 1930. This report is of particular interest in so far as it is the first report issued after the staff reorganisation. The Hospital is now run on the permanent unit system which is detailed on another page. Several additions have been made to the visiting obstetrical staff. Two visiting anæsthetists have been appointed, and a senior resident obstetric officer has been appointed.

The Hospital, which is the largest purely maternity hospital in the country, has 175 beds. Of these, 78 are reserved for Ante-natal cases, 78 for lying-in cases, and 19 are set aside for "suspect" cases. These latter beds are situated in a separate block, the isolation block. Any cases of puerperal sepsis which arise are transferred to the Corporation Fever Hospitals by arrangement with the Public Health authorities.

The Maternity Hospital has continued its policy of not refusing treatment to any woman in labour, but owing to the increasing demands upon the Hospital in recent years, this has only been made possible by co-operation with the Public Health authorities, who have accommodated the overflow cases (usually normal cases) in their hospitals. These "overflow" cases are dealt with in an Appendix and are not considered in making up the statistical report which deals only with cases actually treated in the Maternity Hospital.

A new admission room has been added to the Hospital embracing many modern improvements, including sprays, baths, and an emergency labour room.

An additional labour ward, with separate first stage rooms and rest rooms, etc., has been built and should prove of great service to the Hospital.

The Ante-natal Department, which is becoming an increasingly important part of the Hospital, carries on its work in co-operation with the Ante-natal Dispensary (on the premises), from which many cases are admitted, either to the Ante-natal wards or to the Obstetrical Department.

The Ante-natal Dispensary is yearly becoming more popular, and the Directors have found it necessary to open it on Saturdays, thus affording facilities for attendance on all days except Sundays. With

this Ante-natal Dispensary are combined a Post-natal Dispensary and an Infant Consultation or Child Welfare Centre, at all of which the total attendances during the year 1930 numbered 19,588.

In addition to the indoor cases and dispensary cases referred to above the Hospital carries on a large amount of obstetrical work in "the district," *i.e.*, in the patients' own homes. During the year 1930, 4,079 women were attended in childbirth by the Maternity staff, and, in all, 44,894 visits were paid to them in their homes.

Many patients are sent into Hospital as emergency cases by medical practitioners and midwives, and even by other institutions. Many cases are sent from a considerable distance, *e.g.*, the Western Isles.

During the year 1930, 70·1 per cent. of the admissions to Hospital were abnormal cases. Many of these were emergency cases. This fact, together with many others, such as the high incidence of rickets in the population which is served by the Hospital, indicate that the type of case dealt with is unfortunately often of the gravest variety. This has to be borne in mind when comparing the incidence of the more serious complications of labour and the mortality and morbidity rates with those of institutions where there is not this measure of abnormality with which to contend.

The report has been drawn up as far as possible along the lines suggested by the Committee of the Royal Society of Medicine for the unification of maternity hospital reports. The patients are divided into three categories. The first, category A, are those which have received some measure of ante-natal care. The second category B, are the emergency cases, the majority of whom have received little or no ante-natal care. The third, category C (not specified by the Committee of the Royal Society of Medicine), are those patients who apply at the Hospital for admission without either ante-natal care or doctor's recommendation. These categories are referred to in more detail in the text.

It may be of interest to the reader who is not personally acquainted with the work of the Hospital, to mention that a considerable amount of research work is being carried on by members of the medical staff into urinary infections, puerperal fever, toxæmia of pregnancy, cardiac disease, etc.

During the year 1930 the number of patients treated by the Hospital was 8,548. This number does not include the patients attending the Ante-natal and Post-natal Dispensaries, at which there were 19,588, as has been already mentioned.

Of a total of 9,326 patients, 4,469 were admitted to Hospital, 778 were sent to the Public Health Hospitals, and 4,079 were treated in their own homes in the district. The patients treated in the wards thus numbered 4,469, and it is with these patients that this report deals.

The cases have been divided into categories, the number in each being as follows :—

Category A (cases which had been under ante-natal supervision)	-	-	-	-	-	-	-	1,790
Category B (emergency cases)	-	-	-	-	-	-	-	1,934
Category C (cases applying for admission without recommendation of any kind)	-	-	-	-	-	-	-	745
								<hr/>
Total	-	-	-	-	-	-	-	4,469
								<hr/> <hr/>

From these figures it is seen that of the total of 4,469 patients treated in the wards of the Hospital, 1,790 (or 40 per cent.) had been under ante-natal supervision of some kind, 1,934 (or 43·2 per cent.) were sent in as emergency cases, and 745 (or 16·6 per cent.) were admitted without recommendation of any kind.

The total number of cases treated in the wards of the Ante-natal Department was 1,579 (or 35·3 per cent. of all the indoor cases). The apparent discrepancy between this figure and the total number of cases which had been under ante-natal supervision is due mainly to the direct admission of patients from the Ante-natal Dispensary to the Obstetrical Department.

*Category A—Cases which had been under Ante-natal
Supervision.*

1. Delivered in hospital, -	-	-	-	-	-	-	-	1233
2. Died undelivered, -	-	-	-	-	-	-	-	4
3. B.B.A. (child born before admission),	-	-	-	-	-	-	-	3
4. Abortions, -	-	-	-	-	-	-	-	151
5. Ectopic gestation, -	-	-	-	-	-	-	-	4
6. Miscellaneous (including cases dismissed undelivered),	-	-	-	-	-	-	-	395
Total, -								<hr/> 1790 <hr/>
Died (including 4 who died undelivered),	-	-	-	-	-	-	-	23

Category B—Cases sent in as Emergencies.

1. Delivered in hospital -	-	-	-	-	-	-	-	1016
2. Died undelivered, -	-	-	-	-	-	-	-	16
3. B.B.A. (child born before admission),	-	-	-	-	-	-	-	25
4. Abortions, -	-	-	-	-	-	-	-	592
5. Ectopic gestation, -	-	-	-	-	-	-	-	4
6. Miscellaneous (including cases dismissed undelivered),	-	-	-	-	-	-	-	281
Total, -								<hr/> 1934 <hr/>
Died (including 16 who died undelivered),	-	-	-	-	-	-	-	85

Category C—Cases not under Ante-natal Supervision.

1. Delivered in hospital, -	-	-	-	-	-	-	-	528
2. Died undelivered, -	-	-	-	-	-	-	-	—
3. B.B.A. (child born before admission),	-	-	-	-	-	-	-	4
4. Abortions, -	-	-	-	-	-	-	-	179
5. Ectopic gestation -	-	-	-	-	-	-	-	3
6. Miscellaneous (including cases dismissed undelivered),	-	-	-	-	-	-	-	31
Total, -								<hr/> 745 <hr/>
Died, -	-	-	-	-	-	-	-	11

NUMERICAL SUMMARY OF CASES DELIVERED IN HOSPITAL OR
ADMITTED AFTER DELIVERY.

Presentations (Categories A, B, and C).

Note.—This table does not include cases which were dismissed undelivered.

Presentation.	Prims.	Multips.	Total.
Vertex—			
Occipito-anterior - - - -	942	1322	2264
Occipito-posterior - - - -	63	48	111
Face (unclassified) - - - -	4	26	30
Total - -	1009	1396	2405
Breech - - - - -	82	111	193
Transverse - - - - -	2	20	22
Presentation not recorded - -	60	97	157
B.B.A. - - - - -	9	23	32
	1162	1647	2809

Category A.—Cases which had been under Ante-natal Supervision.

Excluding cases of abortion and ectopic gestation, 1635 patients were under ante-natal supervision before delivery.

The various abnormalities found in cases which had been under supervision are shown in the following table :—

Abnormality.	Primigravidæ.	Multiparæ.	Total.
Delayed labour—			
Malpresentation or malposition - -	51	84	135
Contracted pelvis - - - -	102	106	208
Normal labour, no delay or abnormality found - - - -	207	409	616
Ante-partum hæmorrhage—			
Placenta prævia - - - -	1	17	18
Accidental hæmorrhage - - - -	3	14	17
Multiple pregnancy—			
Twins - - - - -	4	15	19
Triplets - - - - -	1	—	1
Toxæmia of pregnancy—			
Albuminuria, pre-eclamptic toxæmia, and nephritic toxæmia - - -	62	72	134
Eclampsia - - - - -	4	3	7
Hyperemesis gravidarum - - -	10	22	32
Various foetal states—			
Hydrocephalus - - - - -	2	3	5
Anencephaly - - - - -	1	1	2
Spina bifida - - - - -	—	1	1
General foetal dropsy - - - -	—	1	1
Macerated foetus - - - - -	—	5	5
Various other conditions—			
Vaginal cicatrix - - - - -	—	1	1
Furunculosis - - - - -	—	1	1
General debility - - - - -	—	5	5
Cystic ovary - - - - -	—	1	1
Retroverted gravid uterus - - -	1	12	13
Septic hand - - - - -	—	1	1
Hydramnios - - - - -	4	12	16
Respiratory diseases other than phthisis	3	21	24
Prolapsed cord - - - - -	—	3	3
Chorea - - - - -	2	5	7
Cystitis - - - - -	1	1	2
Anæmia - - - - -	—	2	2
Bartholin's cyst - - - - -	—	1	1
Trauma - - - - -	—	1	1
Renal calculus - - - - -	—	1	1
Varicose veins - - - - -	—	5	5
Phthisis - - - - -	1	1	2
Fibromyoma - - - - -	1	1	2
Cervical erosion - - - - -	—	1	1
Influenza - - - - -	—	1	1
Hysteria - - - - -	—	1	1
Amputated cervix - - - - -	—	1	1
Typhoid fever - - - - -	—	1	1
Gastric ulcer - - - - -	—	1	1
Epilepsy - - - - -	2	—	2
Pendulous abdomen - - - - -	1	—	1
Salpingitis - - - - -	—	1	1
Vaginal cyst - - - - -	—	1	1
Endometritis - - - - -	1	—	1
Congenital abnormality of uterus - -	2	—	2
Lactosuria - - - - -	—	1	1

Abnormalities found in Cases—continued.

Abnormality.	Primigravidæ.	Multiparæ.	Total.
Lumbago - - - - -	—	1	1
Cerebral syphilis - - - - -	—	1	1
Exophthalmic goitre - - - - -	1	—	1
Hæmatemesis - - - - -	1	2	3
Prolapse of uterus - - - - -	—	2	2
False labour - - - - -	4	12	16
Observation - - - - -	1	5	6
Retained placenta - - - - -	—	3	3
? Pregnancy - - - - -	1	2	3

Category B—Cases sent in as Emergencies.

The number of cases sent in as emergencies was 1,934. This total includes 592 cases of abortion and 4 cases of ectopic gestation or other complication of the earlier months of pregnancy. The remaining cases are classified in the following numerical summary :—

Table showing the incidence of various Emergencies.

Emergency.	Primigravidæ.	Multiparæ.	Total.
Delayed labour—			
Malpresentation or malposition - -	89	112	201
Contracted pelvis - - - - -	62	41	103
Multiple pregnancy—			
Twins - - - - -	6	18	24
Triplets - - - - -	1	—	1
Normal labour : no delay or abnormality found - - - - -	76	126	202
Ante-partum hæmorrhage—			
Placenta prævia - - - - -	8	56	64
Accidental hæmorrhage - - - - -	5	77	82
Toxæmia of pregnancy—			
Albuminuria, pre-eclamptic toxæmia, and nephritic toxæmia - - - - -	51	75	126
Eclampsia - - - - -	34	15	49
Hyperemesis - - - - -	28	47	75
Prolapsed cord - - - - -	8	14	22
? Pregnancy - - - - -	—	3	3
Retroverted gravid uterus - - - - -	1	8	9
Adherent placenta - - - - -	2	1	3
Perineal repair - - - - -	3	4	7
False labour - - - - -	—	5	5
Prolapse of uterus - - - - -	—	1	1
Post-partum hæmorrhage - - - - -	1	4	5
Retained placenta - - - - -	9	18	27
Various foetal states—			
Impacted shoulders - - - - -	—	1	1
Hydrocephalus - - - - -	1	6	7
Anencephaly - - - - -	1	6	7
Hydrocephalus and meningocele - - - - -	1	1	2
Macerated foetus - - - - -	2	1	3

Incidence of various Emergencies—continued.

Emergency.	Primigravidæ.	Multiparæ.	Total.
Various other conditions—			
Respiratory diseases other than			
phthisis - - - - -	3	15	18
Vaginal cicatrix - - - - -	—	1	1
Encephalitis lethargica - - - - -	1	—	1
Carcinoma of liver - - - - -	—	2	2
Cerebral tumour - - - - -	—	1	1
Pseudocyesis - - - - -	—	1	1
Bell's paralysis - - - - -	1	—	1
Cystitis - - - - -	1	—	1
Fibromyoma - - - - -	1	—	1
Anæmia - - - - -	—	2	2
Ischio-rectal abscess - - - - -	—	1	1
Chorea - - - - -	1	1	2
Septic meningitis - - - - -	1	—	1
Mucous polypus - - - - -	—	1	1
Varicose veins - - - - -	—	1	1
Post maturity - - - - -	—	1	1
Jaundice - - - - -	—	1	1
Adhesions - - - - -	—	1	1
Ventral hernia - - - - -	—	1	1
Syncopal attack - - - - -	—	2	2
Phthisis - - - - -	1	1	2
Hæmatemesis - - - - -	—	4	4
Hydramnios - - - - -	—	5	5
Rigidity of cervix - - - - -	1	2	3
Hemiplegia - - - - -	—	1	1
Influenza - - - - -	—	1	1
Epilepsy - - - - -	1	4	5
Bursitis of knee - - - - -	—	1	1
Atresia of cervix - - - - -	—	1	1
Hysteria - - - - -	2	1	3
Colic - - - - -	—	1	1
Ovarian cyst - - - - -	—	1	1
Diabetes - - - - -	—	1	1
Salpingitis - - - - -	—	1	1
Appendicitis - - - - -	—	1	1
Subinvolution - - - - -	—	1	1
Lysol poisoning - - - - -	—	1	1
Anæmia - - - - -	—	2	2
Hydronephrosis - - - - -	—	1	1
Trauma - - - - -	—	2	2
Inversion of uterus - - - - -	—	1	1
Sciatica - - - - -	—	1	1
Septic leg - - - - -	—	1	1
Endometritis - - - - -	—	1	1
Ulcerative colitis - - - - -	—	1	1
Cholecystitis - - - - -	—	1	1
Observation - - - - -	3	6	9
Cystic ovary - - - - -	—	1	1
Debility - - - - -	—	1	1
Anuria - - - - -	—	1	1
Renal colic - - - - -	—	1	1

Category C—Cases applying for admission without Previous Supervision.

The number of cases applying for admission without previous supervision was 745. These cases are classified in the following numerical summary:—

Table showing the incidence of various abnormalities in cases applying for admission without Previous Supervision.

Abnormality.	Primigravidæ.	Multiparæ.	Total.
Delayed labour—			
Malpresentation or malposition - -	17	18	35
Contracted pelvis - - - -	14	8	22
Multiple pregnancy (twins) - - -	3	10	13
Normal labour: no delay or abnormality found - - - - -	159	287	446
Ante-partum hæmorrhage—			
Placenta prævia - - - - -	1	11	12
Accidental hæmorrhage - - - -	2	13	15
Toxæmia of pregnancy—			
Albuminuria, pre-eclamptic toxæmia, and nephritic toxæmia - - -	9	10	19
Eclampsia - - - - -	3	—	3
Hyperemesis - - - - -	—	3	3
Prolapsed cord - - - - -	—	3	3
False labour - - - - -	1	3	4
? Pregnancy - - - - -	1	3	4
Retroverted gravid uterus - - -	—	3	3
Various foetal states—			
Hydrocephalus - - - - -	1	5	6
Anencephaly - - - - -	—	2	2
Macerated foetus - - - - -	—	1	1
Spina bifida - - - - -	—	1	1
Various other conditions—			
Chorea - - - - -	2	—	2
General debility - - - - -	—	2	2
Influenza - - - - -	—	2	2
Hæmorrhoids - - - - -	1	1	2
Varicose veins - - - - -	—	2	2
Pleurisy - - - - -	—	1	1
Anæmia - - - - -	—	2	2
Trauma - - - - -	—	2	2
Ventral hernia - - - - -	—	1	1
Salpingo oophoritis - - - - -	—	1	1
Hydramnios - - - - -	1	6	7
Œdema of vulva - - - - -	—	1	1
Encephalitis lethargica - - - -	—	1	1
Medico legal cause - - - - -	1	—	1
Subinvolution - - - - -	—	1	1
Cystitis - - - - -	—	1	1
Salpingitis - - - - -	—	1	1
Respiratory diseases other than phthisis - - - -	—	2	2
Varicose veins - - - - -	—	1	1
Enteritis - - - - -	—	1	1
Complete perineal repair - - -	—	1	1

ANTE-NATAL CASES—INDOOR.

There were 1,571 cases treated in the wards of the Ante-natal Department during the year. This total, equivalent to 35.1 per cent. of all indoor cases, is to be distinguished from the total of 1,790 cases which were under ante-natal supervision (see page 7). The following table shows the disease or accident of pregnancy for which treatment was undertaken in the Ante-natal Wards, and the results :—

Cases admitted to Ante-natal Wards : Disease or Accident of Pregnancy for which treatment was undertaken.

Disease or Accident of Pregnancy.	No. of Cases.	Mother.	
		Lived.	Died.
1. Toxæmia of pregnancy—			
(a) Albuminuria, pre-eclamptic toxæmia and nephritic toxæmia - -	312	309	3
(b) Eclampsia - - - - -	18	17	1
(c) Hyperemesis - - - - -	144	142	2
2. Ante-partum hæmorrhage - - - - -	93	91	2
3. Threatened abortion - - - - -	263	262	1
4. Pulmonary disease - - - - -	34	32	2
5. Heart disease - - - - -	132*	130	2
6. Contracted pelvis - - - - -	200	199	1
7. Phthisis - - - - -	3	3	—
8. Other obstetrical complications - -	144	143	1
9. Various other conditions other than above - - - - -	228	225	3
Totals - -	1,571	1,553	18

* Includes re-admissions.

OBSTETRICAL CASES—INDOOR.

MALPRESENTATIONS.

1. *Vertex—Occipito-posterior.*

There were 113 cases in which there was an occipito-posterior presentation. There were two maternal deaths and 27 stillbirths. The following table shows the mode of delivery and the results of these cases :—

Cases of Occipito-posterior (Vertex) Presentation.

Mode of Delivery.	No. of Cases.				Results. Mother lived.				Child born alive.			
	A	B	C	Total.	A	B	C	Total.	A	B	C	Total.
Spontaneous delivery	23	10	6	39	23	10	6	39	23 ¹	9 ²	5 ³	37
Forceps - - -	22	37	2	61	21 ⁴	36 ⁵	2	59	20 ⁶	25 ⁷	1 ⁸	46
Cæsarean section -	1	2	—	3	1	2	—	3	1	2	—	3
Version to breech -	1	—	—	1	1	—	—	1	— ⁹	—	—	—
Craniotomy - -	2	3	1	6	2	3	1	6	—	—	—	10
Undelivered - -	2	—	—	2	2	—	—	2	—	—	—	—
Manual delivery (im- pacted shoulders) -	—	—	1	1	—	—	—	1	— ¹¹	—	—	—
Totals - - -	51	52	10	113	50	51	9	111	35	35	4	86

¹ 1 Died.

² 3 Stillbirths (twins, twice).

³ 1 Stillbirth.

⁴ 1 Maternal death.

⁵ 1 Maternal death.

⁶ 2 Stillbirths.

⁷ 12 Stillbirths.

⁸ 1 Stillbirth.

⁹ 1 Stillbirth.

¹⁰ 6 Stillbirths.

¹¹ 1 Stillbirth.

2. Breech.

There were 205 cases of breech presentation ; 77 were in patients who had been under ante-natal supervision ; 106 were emergency cases ; 22 were in patients who applied for admission without having been under previous supervision. There were 6 maternal deaths and 49 stillbirths. Since primiparity in itself constitutes a complication of a breech case, primigravidæ have been kept in a separate group in the table which follows :—

Breech Presentation.

(a) Primigravidæ.

Mode of Delivery.	No. of Cases.				Results. Mother lived.				Child born alive.			
	A	B	C	Total.	A	B	C	Total.	A	B	C	Total.
Spontaneous delivery	3	5	1	9	3	5	1	9	3 ¹	6 ²	1	10
Manual delivery, manipulation of limbs - - -	14	40	10	64	14	39 ³	10	63	11 ⁴	30 ⁵	7 ⁶	48
Forceps - - -	1	6	—	7	1	6	—	7	— ⁷	3 ⁸	—	3
Cæsarean section -	1	—	—	1	1	—	—	1	1	—	—	1
Version, spontaneous delivery - - -	—	—	1	1	—	—	1	1	—	—	— ⁹	—
Undelivered - - -	4	—	—	4	4	—	—	4	—	—	—	—
Totals - - -	23	51	12	86	23	50	12	85	15	39	8	62

¹ 1 Stillbirth (twins, once).

² Twins (once).

³ 1 Maternal death.

⁴ 4 Stillbirths (twins, once).

⁵ 11 Stillbirths (twins, once).

⁶ 3 Stillbirths.

⁷ 1 Stillbirth.

⁸ 3 Stillbirths.

⁹ 1 Stillbirth.

(b) *Multiparæ.*

Mode of Delivery.	No. of Cases.				Results. Mother lived.				Child born alive.			
	A	B	C	Total.	A	B	C	Total.	A	B	C	Total.
Spontaneous delivery	34	18	4	56	34	17 ¹	4	55	34 ²	14 ³	5 ⁴	53
Manual delivery, manipulation of limbs - - -	12	33	6	51	11 ⁵	31 ⁶	5 ⁷	47	10 ⁸	22 ⁹	6 ¹⁰	38.
Version to vertex -	1	1	—	2	1	1	—	2	— ¹¹	1	—	1
Perforation of head -	—	2	—	2	—	2	—	2	—	— ¹²	—	—
Undelivered - -	7	1	—	8	7	1	—	8	—	—	—	—
Totals - - -	54	55	10	119	53	52	9	114	44	37	11	92

- ¹ 1 Maternal death.

² 2 Stillbirths (twins, twice).

³ 5 Stillbirths (twins, once).

⁴ 1 Stillbirth. (twins, twice).

⁵ 1 Maternal death

⁶ 2 Maternal deaths.
- ⁷ 1 Maternal death.

⁸ 2 Stillbirths.

⁹ 11 Stillbirths.

¹⁰ 1 Stillbirth (twins, once).

¹¹ 1 Stillbirth.

¹² 2 Stillbirths.

(c) *Primigravidæ and Multiparæ.*

Mode of Delivery.	No. of Cases.				Results. Mother lived.				Child born alive.			
	A	B	C	Total.	A	B	C	Total.	A	B	C	Total.
Spontaneous delivery	38	24	6	68	38	23	6	67	37	20	5	62
Manual delivery, manipulation of limbs - - -	26	73	16	115	25	70	15	110	23	52	13	88
Forceps - - -	1	6	—	7	1	6	—	7	—	3	—	3
Cæsarean section -	1	—	—	1	1	—	—	1	1	—	—	1
Perforation of head -	—	2	—	2	—	2	—	2	—	—	—	—
Undelivered - -	11	1	—	12	11	1	—	12	—	—	—	—
Totals - - -	77	106	22	205	76	102	21	199	61	75	18	154

Maternal deaths, 6 (= 2·9 per cent.) Stillbirths, 49 (= 25·3 per cent.) Twins, 10 sets.

3. *Face and Brow.*

There were 30 cases of face or brow presentation. There was one maternal death and 10 stillbirths. The following table shows the mode of delivery and the results in these cases:—

Face and Brow (Vertex) Presentation.

Mode of Delivery.	No. of Cases.				Results. Mother lived.				Child born alive.			
	A	B	C	Total.	A	B	C	Total.	A	B	C	Total.
Spontaneous delivery	3	10	2	15	3	10	2	15	3	9 ¹	2	14
Conversion to vertex	—	2	1	3	—	2	1	3	—	2	1	3
Conversion to breech	—	1	—	1	—	1	—	1	—	— ²	—	—
Craniotomy - - -	—	3	—	3	—	2 ³	—	2	—	— ⁴	—	—
Version, decapitation	—	1	—	1	—	1	—	1	—	— ⁵	—	—
Forceps - - - -	1	5	1	7	1	5	1	7	1	1 ⁶	1	3
Totals - - -	4	22	4	30	4	21	4	29	4	12	4	20

¹ 1 Stillbirth (and 1 died).

² 1 Stillbirth.

³ 1 Maternal death.

⁴ 3 Stillbirths.

⁵ 1 Stillbirth.

⁶ 4 Stillbirths.

4. *Transverse.*

There were 23 cases of transverse presentation. There were no maternal deaths and 11 stillbirths. The mode of treatment and the results of these cases are shown in the following table:—

Transverse Presentation.

Mode of Delivery.	No. of Cases.				Results. Mother lived.				Child born alive.			
	A	B	C	Total.	A	B	C	Total.	A	B	C	Total.
Conversion to breech	3	9	—	12	3	9	—	12	3	4 ¹	—	7
Manual - - -	2	—	1	3	2	—	1	3	1 ²	—	— ³	1
Cæsarean section -	—	2	—	2	—	2	—	2	—	2	—	2
Forceps - - -	—	1	—	1	—	1	—	1	—	— ⁴	—	—
Spontaneous - -	—	2	—	2	—	2	—	2	—	1 ⁵	—	1
Perforation of head, blunt hook - -	—	1	—	1	—	1	—	1	—	— ⁶	—	—
Undelivered - -	—	—	1	1	—	—	1	1	—	—	—	—
Craniotomy - -	1	—	—	1	1	—	—	1	—	— ⁷	—	—
Totals - - -	6	15	2	23	6	15	2	23	4	7	—	11

¹ 5 Stillbirths (and 1 died).

² 1 Stillbirth.

³ 1 Stillbirth.

⁴ 1 Stillbirth.

⁵ 1 Stillbirth.

⁶ 1 Stillbirth.

⁷ 1 Stillbirth.

MULTIPLE PREGNANCY—1. *Twins.*

There were twins in 56 cases during the year, 13 in primigravidæ and 43 in multiparæ. The following tables summarise the main points:—

Category A—Cases which had been under Ante-natal Supervision.

Reg. No.	Presentation.		Sexes.		Premature.	Mature.	Weights.		Results.			Remarks.
	1	2	1	2			1	2	M.	Ch. I.	Ch. II.	
94	Vertex	Breech	F	F	+	—	4½	7	A	A	A	Hydramnios. Forceps.
146	Breech	Breech	F	F	—	+	4¾	4	A	A	A	
495	Breech	Breech	F	M	—	+	7½	7½	A	A	A	
780	Vertex	Breech	F	M	—	+	5¼	6	D	A	S.B.	
794	Vertex	Breech	M	M	—	+	6½	6	A	A	A	Forceps & Manual delivery C. P. Albuminuria.
803	Breech	Breech	F	M	—	+	4½	4½	A	A	A	
1069	Vertex	Breech	F	F	+	—	2½	2¼	A	A	A	
1107	Vertex	Vertex	F	M	+	—	3	3	A	A	A	
1189	Vertex	Breech	F	F	—	+	7	5	A	A	A	Forceps. Forceps.
1832	—	—	F	M	—	+	5¼	6	A	A	A	
2140	Vertex	Vertex	F	F	—	+	7¼	7	A	A	A	
2505	Vertex	Vertex	M	F	—	+	6¾	6	A	A	A	
2925	Breech	Breech	F	M	+	—	5¼	5	A	A	A	Forceps.
3517	Tr.	Breech	F	F	—	+	5½	5¼	A	A	A	
3801	Breech	Breech	F	F	+	—	3½	4½	A	S.B.	S.B.	
3865	Vertex	Breech	M	M	+	—	3½	3½	A	A	A	
3862	Vertex	Breech	M	M	—	+	6¾	8	A	A	A	Forceps.
4197	Vertex	Vertex	F	F	+	—	4¼	4½	A	A	A	
4148	Breech	Breech	M	M	+	—	5¾	4¾	A	A	A	

Category B—Cases sent in as Emergencies.

Reg. No.	Presentation.		Sexes.		Premature.	Mature.	Weights.		Results.			Remarks.
	1	2	1	2			M.	Ch. I.	Ch. II.			
500	Breech	Vertex	F	M	—	+	6	6	A	A	A	Placenta prævia. Albuminuria.
578	Breech	Breech	M	M	+	—	5½	5	A	A	A	
767	Vertex	Vertex	M	M	—	+	3	3¼	D	A	A	
1060	Vertex	Breech	F	F	+	—	4¼	4¾	A	S.B.	A	
1315	Vertex	Brow	M	M	+	—	3½	4½	A	A	S.B.	F.F.O. Forceps.
1360	Vertex	Breech	M	M	—	+	8½	6¼	D	S.B.	S.B.	
1572	—	—	M	M	+	—	5	4½	D	S.B.	S.B.	
1607	Face	Breech	M	M	+	—	1¾	2½	A	A	A	
1654	Vertex	Vertex	F	F	—	+	4½	6	A	A	S.B.	Forceps.
1720	Vertex	Breech	F	F	—	+	5	5	A	A	A	
1934	Breech	Breech	F	F	—	+	7½	7¼	A	A	A	
1986	Vertex	Breech	F	F	—	+	5	5¼	A	A	A	
2074	Vertex	Vertex	M	F	—	+	6¾	7	A	A	A	Albuminuria.
2205	Vertex	Breech	M	M	—	+	6¾	6¼	D	S.B.	S.B.	
2241	Breech	Breech	M	F	—	+	8¼	—	A	S.B.	A	
2245	Breech	Vertex	M	F	—	+	6	4¾	A	S.B.	A	
2422	Vertex	Breech	M	F	—	+	7	6½	A	A	A	Albuminuria.
2572	Breech	Vertex	M	F	+	—	3¼	3	A	A	A	
2837	Breech	Breech	M	F	+	—	5¾	5	A	A	A	
2915	Breech	Vertex	M	F	+	—	5¼	6	A	A	A	
3132	Vertex	Breech	F	M	—	+	5	6¼	A	A	A	Acc. hæm. Albuminuria.
4047	Vertex	Vertex	F	F	+	—	3¼	2¾	A	S.B.	S.B.	
4158	Breech	Breech	M	F	+	—	5¼	5½	A	A	A	
4383	Vertex	Breech	M	F	+	—	1½	3	A	S.B.	A	

Category C—Cases not under Ante-natal Supervision.

Reg. No.	Presentation.		Sexes.		Premature.	Mature.	Weights.		Results.			Remarks.
	1	2	1	2			1	2	M.	Ch. I.	Ch. II.	
390	Vertex	Vertex	F	F	+	—	3 $\frac{1}{4}$	3	A	A	S.B.	Placenta prævia. Album-inuria.
821	Breech	Vertex	F	F	—	+	5 $\frac{1}{2}$	5	A	A	A	
1128	—	—	M	M	+	—	4 $\frac{1}{2}$	4 $\frac{1}{2}$	A	A	A	
1478	Vertex	Vertex	F	F	—	+	5 $\frac{3}{4}$	4 $\frac{1}{2}$	A	A	A	
1662	Vertex	Vertex	M	M	—	+	6 $\frac{3}{4}$	6 $\frac{1}{2}$	A	A	A	
2286	Vertex	Breech	M	M	—	+	5 $\frac{1}{4}$	5	A	A	A	
2784	Vertex	Vertex	M	M	+	—	5	5 $\frac{1}{4}$	A	A	A	
2933	Breech	Face	M	F	—	+	6 $\frac{1}{4}$	6	A	A	A	
3053	Vertex	Breech	M	F	—	+	5 $\frac{1}{2}$	6	D	A	S.B.	
3342	Vertex	Vertex	M	M	—	+	4 $\frac{1}{2}$	3 $\frac{3}{4}$	A	A	A	
3538	Vertex	Breech	F	M	+	—	3 $\frac{1}{4}$	3 $\frac{3}{4}$	A	A	A	
3863	Vertex	Vertex	F	F	+	—	2 $\frac{3}{4}$	3 $\frac{1}{4}$	A	A	A	
3897	Breech	Vertex	F	F	+	—	6 $\frac{1}{4}$	4 $\frac{3}{4}$	A	A	A	

2. Triplets.

Reg. No.	Presentation.			Sexes.			Premature.	Mature.	Weights.			Results.				Remarks.
	1	2	3	1	2	3			1	2	3	M.	Ch. I.	Ch. II.	Ch. III.	
1340	V.	Br.	Br.	M	M	F	—	+	5	5	5½	A	A	A	A	Forceps. 1 child died.
2811	V.	Br.	Br.	F	F	M	—	+	4½	3¾	3¾	A	A	A	A	

OBSTETRICAL OPERATIONS.

The number of obstetric operations performed during the year 1930 was 645. A summary of these cases is given in the following tables :—

1. *Forceps.*

Forceps were applied in 297 cases, and the results are shown in the following table :—

Indication.	No. of Cases.	Mother.		Child.		
		Alive.	Dead.	Alive.	S.B.	Died.
Contracted pelvis - - -	69	67	2	52	14	3
Dystocia from various causes -	100	97	3	78 ¹	19	5
Toxæmia of pregnancy - -	39	36	3	28 ²	10	2
Failed forceps outside - -	25	21	4	12	9	4
Prolapsed cord - - - -	2	2	—	—	2	—
Ante-partum hæmorrhage - -	5	4	1	—	4	1
Malposition or malpresentation -	48	47	1	32 ³	16	1
Heart disease - - - -	9	9	—	9	—	—
Totals - - - -	297	283	14	211	74	16

¹ Twins (twice).² Twins.³ Twins.

2. Induction.

Surgical induction was performed in 63 cases, and the results are shown in the following table :—

Indication.	No. of Cases.	Mother.		Child.			
		Alive.	Died.	Alive.	S.B.	Died.	Misc.
Toxæmia of pregnancy -	47	45	2	14 ¹	10 ²	8	15
Heart disease - -	1	1	—	—	1	—	—
Inevitable abortion -	4	4	—	—	—	—	4
Chorea - - - -	2	1	1	—	—	1 ³	—
Hydramnios - - -	1	1	—	—	—	—	1
Pyelitis - - - -	6	4	2	—	1	2	3
Post-maturity - -	1	1	—	1	—	—	—
Contracted pelvis -	1	1	—	1	—	—	—
Totals -	63	58	5	16	12	11	23

¹ 1 Case undelivered. ² 1 Case of twins. ³ 1 Case undelivered.

3. Cæsarean Section.

Cæsarean section was performed in 184 cases, and the results are shown in the following table :—

Indication.	No. of Cases.	Mother.		Child.		
		Alive.	Died.	Alive.	S.B.	Died.
Contracted pelvis - - -	121	114	7	117	4	—
Ante-partum hæmorrhage - -	33	32	1	29	3	1
Heart disease - - - -	21	18	3	20	—	1
Elderly primipara - - - -	3	3	—	3	—	—
Cervical carcinoma - - - -	1	—	1	1	—	—
Atresia of cervix - - - -	1	1	—	1	—	—
Chorea - - - - -	1	1	—	—	1	—
Ruptured uterus - - - -	1	—	1	—	1	—
Habitual stillbirths - - - -	1	1	—	1	—	—
Spondylolisthesis - - - -	1	1	—	1	—	—
Totals - - -	184	171	13	173	9	2

Maternal death-rate = 7·0 per cent.
Infantile death-rate = 5·4 per cent.

Cæsarean Section—A.

Reg. No.	Category.	Age.	Parity.	Stage of Pregnancy.	Indication for Operation.	Mother.		Child.		
						Alive.	Died.	Alive.	S.B.	Died.
3	A	41	1	F T	Elderly primipara	+	—	+	—	—
61	A	35	10	F T	Cervical carcinoma	—	+	+	—	—
207	A	21	1	F T	Contracted pelvis	+	—	+	—	—
208	A	25	1	F T	C.P.	+	—	+	—	—
247	A	37	4	F T	C.P.	+	—	+	—	—
359	A	21	1	F T	Contracted pelvis	+	—	+	—	—
408	A	48	3	F T	Habitual still births	+	—	+	—	—
514	A	25	2	F T	Contracted pelvis	+	—	+	—	—
640	A	—	1	F T	Contracted pelvis	+	—	+	—	—
764	A	38	4	—	Cardiac	—	+	—	+	—
785	A	27	3	F T	C.P.	+	—	+	—	—
811	A	26	1	F T	C.P.	+	—	+	—	—
832	A	—	—	—	Cardiac	+	—	+	—	—
882	A	20	1	F T	Contracted pelvis	+	—	+	—	—
913	A	39	1	F T	Elderly primipara	+	—	+	—	—
977	A	26	3	F T	C.P.	+	—	+	—	—
1005	A	23	1	F T	C.P.	+	—	+	+	—
1007	A	26	1	F T	Contracted pelvis	+	—	+	—	—
1129	A	27	2	F T	C.P.	+	—	+	—	—
1141	A	21	1	F T	C.P.	+	—	+	—	—
1146	A	26	2	F T	C.P.	+	—	+	—	—
1158	A	36	2	P	Placenta prævia	+	—	—	+	—
1170	A	31	3	F T	C.P.	+	—	+	—	—
1172	A	29	1	F T	C.P.	+	—	+	—	—
1232	A	25	1	F T	C.P.	+	—	+	—	—
1234	A	27	4	F T	C.P.	—	+	+	—	—
1243	A	27	2	F T	C.P.	+	—	+	—	—
1252	A	40	2	F T	C.P.	+	—	+	—	—
1259	A	30	7	F T	Placenta prævia	+	—	+	—	—
1287	A	37	2	F T	C.P.	+	—	+	—	—
1327	A	35	4	F T	C.P.	+	—	+	—	—
1383	A	26	2	F T	C.P.	+	—	+	—	—
1453	A	38	1	F T	C.P.	+	—	+	—	—
1459	A	30	1	F T	C.P.	+	—	+	—	—
1557	A	31	6	F T	C.P.	+	—	+	—	—
1565	A	28	1	F T	C.P.	+	—	+	—	—
1566	A	25	3	F T	C.P.	+	—	+	—	—
1580	A	36	1	F T	Cardiac	+	—	+	—	—
1628	A	25	1	F T	C.P.	+	—	+	—	—
1647	A	38	8	F T	Cardiac	+	—	+	—	—
1653	A	29	4	P	C.P.	+	—	+	—	—
1655	A	23	3	F T	C.P.	+	—	+	—	—
1832	A	30	4	F T	C.P.	+	—	+	—	—
1897	A	26	3	F T	A.P.H.	+	—	+	—	—
1911	A	—	4	F T	Cardiac	+	—	+	—	—
1924	A	29	2	F T	C.P.	+	—	+	—	—
1959	A	35	3	F T	C.P.	+	—	+	—	—
2005	A	22	1	F T	C.P.	+	—	+	—	—
2039	A	24	2	F T	C.P.	+	—	+	—	—

Reg. No.	Category.	Age.	Parity.	Stage of Pregnancy.	Indication for Operation.	Mother.		Child.		
						Alive.	Died.	Alive.	S.B.	Died.
2068	A	36	2	F T	C.P.	+	-	+	-	-
2097	A	36	5	P	Cardiac	+	-	+	-	-
2151	A	18	1	F T	C.P.	+	-	+	-	-
2158	A	40	13	F T	Cardiac	+	-	+	-	-
2170	A	24	3	F T	C.P.	+	-	+	-	-
2191	A	22	1	F T	C.P.	+	-	+	-	-
2196	A	26	8	F T	Placenta prævia	+	-	+	-	-
2237	A	22	1	F T	C.P.	+	-	+	-	-
2239	A	30	5	F T	C.P.	+	-	+	-	-
2316	A	27	5	F T	C.P.	+	-	+	-	-
2350	A	25	3	F T	C.P.	+	-	+	-	-
2367	A	30	4	F T	C.P.	+	-	+	-	-
2579	A	28	4	P	Cardiac	+	-	+	-	-
2692	A	21	3	F T	Placenta prævia	+	-	+	-	-
2710	A	23	1	F T	C.P.	+	-	+	-	-
2729	A	33	2	F T	C.P.	+	-	+	-	-
2777	A	24	1	F T	C.P.	+	-	+	-	-
2788	A	24	1	F T	C.P.	+	-	+	-	-
2853	A	32	5	F T	Cardiac	+	-	+	-	-
2858	A	24	1	P	C.P.	+	-	+	-	-
2899	A	36	7	F T	Cardiac	+	-	+	-	-
2935	A	28	2	F T	Cardiac	+	-	+	-	-
2999	A	39	2	F T	C.P.	+	-	+	-	-
3009	A	33	6	F T	Cardiac	-	+	+	-	-
3048	A	24	2	F T	C.P.	+	-	+	-	-
3052	A	24	3	F T	C.P.	+	-	+	-	-
3204	A	30	5	P	C.P.	+	-	+	-	-
3268	A	30	2	F T	C.P.	+	-	+	-	-
3280	A	25	3	F T	C.P.	-	+	+	-	-
3290	A	27	1	F T	C.P.	+	-	+	-	-
3353	A	32	4	F T	C.P.	+	-	+	-	-
3408	A	32	4	F T	C.P.	+	-	+	-	-
3421	A	—	6	F T	C.P.	+	-	+	-	-
3426	A	31	5	F T	C.P.	+	-	+	-	-
3481	A	18	1	F T	C.P.	+	-	+	-	-
3493	A	23	2	P	Cardiac	+	-	+	-	-
3494	A	27	2	P	Cardiac	+	-	+	-	-
3514	A	21	1	F T	C.P.	+	-	+	-	-
3598	A	24	1	F T	C.P.	+	-	+	-	-
3613	A	25	1	F T	C.P.	+	-	+	-	-
3680	A	28	2	F T	C.P.	+	-	+	-	-
3681	A	22	3	F T	C.P.	+	-	+	-	-
3682	A	28	2	F T	C.P.	+	-	+	-	-
3690	A	27	2	F T	C.P.	+	-	+	-	-
3699	A	19	1	F T	C.P.	+	-	+	-	-
3714	A	21	1	F T	C.P.	+	-	+	-	-
3841	A	35	1	F T	C.P.	+	-	+	-	-
3883	A	24	2	F T	C.P.	+	-	+	-	-
3896	A	29	2	F T	C.P.	+	-	+	-	-
3903	A	17	1	F T	C.P. Cardiac	+	-	+	-	-
3993	A	26	1	F T	C.P.	+	-	+	-	-
4061	A	21	1	F T	G.C.P.	+	-	+	-	-

Reg. No.	Category.	Age.	Parity.	Stage of Pregnancy.	Indication for Operation.	Mother.		Child.		
						Alive.	Died.	Alive	S.B.	Dead.
4081	A	29	2	P	C.P.	+	—	+	—	—
4136	A	24	2	F T	C.P.	+	—	+	—	—
4167	A	25	2	F T	C.P.	+	—	+	—	—
4189	A	23	2	F T	C.P.	+	—	+	—	—
4201	A	30	6	F T	C.P.	+	—	+	—	—
4348	A	24	1	F T	C.P.	+	—	+	—	—
4395	A	30	5	F T	C.P.	+	—	+	—	—

Cæsarean Section—B.

48	B	27	2	P	Placenta prævia	+	—	+	—	—
202	B	24	2	F T	Contracted pelvis	+	—	+	—	—
249	B	25	1	F T	C.P.	+	—	+	—	—
420	B	40	12	F T	Ruptured uterus	—	+	—	+	—
542	B	24	1	F T	C.P.	—	+	+	—	—
606	B	23	1	F T	C.P.	+	—	+	—	—
669	B	30	1	F T	Contracted pelvis	+	—	—	+	—
718	B	22	1	F T	Contracted pelvis	+	—	+	—	—
752	B	33	2	P	Placenta prævia	+	—	+	—	—
769	B	—	7	P	Cardiac	—	+	+	—	—
857	—	36	1	F T	Placenta prævia	+	—	+	—	—
881	B	37	1	F T	Contracted pelvis	+	—	+	—	—
930	B	37	3	F T	C.P.	+	—	+	—	—
1017	B	29	3	P	Placenta prævia	+	—	+	—	—
1025	B	43	12	F T	Accidental hæmorrhage	+	—	—	+	—
1156	B	35	4	P	Placenta prævia	+	—	+	—	—
1209	B	26	5	F T	Cardiac	+	—	+	—	—
1268	B	35	2	F T	C.P.	+	—	+	—	—
1307	B	34	7	F T	Cardiac	+	—	+	—	—
1363	B	33	1	F T	C.P.	+	—	+	—	—
1370	B	27	6	F T	C.P.	+	—	+	—	—
1385	B	20	1	F T	C.P.	+	—	+	—	—
1599	B	38	8	F T	Placenta prævia	+	—	+	—	—
1606	B	30	1	F T	C.P.	+	—	+	—	—
1623	B	30	1	F T	Placenta prævia	+	—	+	—	—
1665	B	25	2	F T	C.P.	+	—	+	—	—
1683	B	29	3	F T	Cardiac	+	—	+	—	—
1807	B	36	8	P	Atresia of cervix	+	—	+	—	—
1808	B	34	2	F T	Placenta prævia	+	—	+	—	—
1848	B	32	2	P	Placenta prævia	+	—	+	—	—
1899	B	27	1	F T	C.P.	+	—	+	—	—
2049	B	25	2	F T	C.P.	+	—	+	—	—
2050	B	28	5	F T	Placenta prævia	+	—	+	—	—
2188	B	40	12	—	Placenta prævia	+	—	+	—	—
2190	B	39	6	F T	Placenta prævia	+	—	+	—	—
2321	B	34	4	F T	Spondylolisthesis	+	—	+	—	—
2482	B	26	1	F T	C.P.	+	—	+	—	—
2515	B	27	4	—	Chorea	+	—	—	+	—
2600	B	29	3	F T	Placenta prævia	+	—	+	—	—
2612	B	36	9	F T	Placenta prævia	+	—	+	—	—
2679	B	36	6	F T	C.P.	+	—	+	—	—

Reg. No.	Category.	Age.	Parity.	Stage of Pregnancy.	Indication for Operation.	Mother.		Child.		
						Alive.	Died.	Alive.	S.B.	Died.
2747	B	32	3	P	Placenta prævia	+	-	+	-	-
2800	B	34	1	P	Placenta prævia	+	-	+	-	-
2817	B	33	4	P	Placenta prævia	+	-	+	-	-
2852	B	33	4	F T	Cardiac	+	-	+	-	+
2874	B	31	1	F T	C.P.	+	-	+	-	-
2969	B	25	1	F T	F.F.O.	+	-	+	-	-
3101	B	25	1	F T	C.P.	-	+	+	-	-
3137	B	43	10	P	Placenta prævia	+	-	+	-	-
3239	B	36	2	P	Placenta prævia	+	-	+	-	-
3241	B	40	5	F T	Placenta prævia	-	+	+	-	-
3292	B	25	1	F T	C.P.	-	+	+	-	-
3449	B	32	3	F T	Placenta prævia	+	-	+	-	-
3498	B	43	1	F T	Elderly primipara	+	-	+	-	-
3537	B	34	10	P	Cardiac	+	-	+	-	-
3749	B	38	6	P	Cardiac	+	-	+	-	-
3805	B	26	4	F T	F.F.O.—C.P.	-	+	-	+	-
3938	B	20	1	F T	C.P.	+	-	+	-	-
3981	B	26	4	F T	C.P.	+	-	+	-	-
4070	B	23	1	F T	C.P.	+	-	+	-	-
4199	B	29	1	F T	Placenta prævia	+	-	+	-	-
4273	B	37	12	F T	A.P.H.	+	-	-	+	-
4295	B	43	4	F T	F.F.O.	+	-	-	+	-
4314	B	37	10	F T	C.P.	-	-	+	-	-

Cæsarean Section—C.

537	C	23	1	F T	Contracted pelvis	+	-	+	-	-
950	C	23	3	F T	Contracted pelvis	+	-	+	-	-
1036	C	40	8	P	Placenta prævia	+	-	+	-	-
1128	C	41	4	P	Placenta prævia	+	-	+	-	-
1938	C	27	3	P	Placenta prævia	+	-	+	-	-
2606	C	30	3	F T	C.P.	+	-	+	-	-
2610	C	30	5	—	Placenta prævia	+	-	+	-	+
2931	C	29	3	P	C.P.	-	+	+	-	-
3171	C	24	2	F T	C.P.	+	-	+	-	-
3459	C	29	2	F T	C.P.	+	-	+	-	-
3647	C	22	1	F T	C.P.	+	-	+	-	-
3795	C	40	2	F T	C.P.	+	-	+	-	-

4. Version.

Version was performed in 49 cases, and the results are shown in the following table :—

Indication.	No. of Cases.	Mother.		Child.			
		Alive.	Dead.	Alive.	S.B.	Died.	Unde-livered.
Malpresentation - - -	27	25	2	10	14 ¹	1	2
Placenta prævia - - -	20	18	2	2	17	1	—
Prolapsed cord - - -	1	1	—	—	—	—	1
Accidental hæmorrhage -	1	1	—	—	1	—	—
Totals - - -	49	45	4	12	32	2	3

¹ One not viable.

5. Craniotomy.

Craniotomy was performed in 34 cases, and the results are shown in the following table :—

Indication	No. of Cases.	Mother.	
		Alive.	Dead.
Contracted pelvis - - -	10	10	—
Malpresentation - - -	6	5	1
Hydrocephalus - - -	5	4	1
Failed forceps outside - -	8	8	—
Breech. Impacted head -	2	2	—
Trial forceps failed - -	3	3	—
Totals - - -	34	32	2

¹ Includes one case of decapitation.

Craniotomy.

Reg. No.	Category.	Age.	Parity.	Stage of Pregnancy.	Indication for Operation.	Mother.		Child.		
						Alive.	Died.	Alive.	S.B.	Died.
25	A	39	4	F T	C.P.	+	-	-	+	-
69	B	23	1	F T	C.P.	+	-	-	+	-
191	A	40	5	F T	C.P.	+	-	-	+	-
343	B	28	3	F T	Hydrocephalus.	+	-	-	+	-
361	B	39	8	F T	Breech. Impacted head	+	-	-	+	-
364	B	39	1	F T	Malpresentation	-	+	-	+	-
555	B	27	1	F T	F.F.O.	+	-	-	+	-
555	B	27	1	F T	F.F.O.	+	-	-	+	-
704	B	36	5	F T	F.F.O.	+	-	-	+	-
783	B	39	11	F T	F.F.O.	+	-	-	+	-
788	A	37	3	F T	Hydrocephalus	+	-	-	+	-
902	B	32	5	F T	F.F.O.	+	-	-	+	-
1098	B	28	1	F T	F.F.O.	+	-	-	+	-
1305	B	35	1	F T	C.P.	+	-	-	+	-
1341	B	33	1	F T	Malpresentation.	+	-	-	+	-
1636	B	23	1	F T	C.P.	+	-	-	+	-
1852	B	30	3	F T	Malpresentation	+	-	-	+	-
1875	A	32	1	F T	F.F.O.	+	-	-	+	-
1945	B	32	4	F T	F.F.O.	+	-	-	+	-
2472	B	26	1	F T	Trial forceps failed	+	-	-	+	-
2645	C	31	2	F T	C.P.	+	-	-	+	-
3003	B	25	4	F T	Malpresentation.	+	-	-	+	-
3438	B	36	1	F T	Hydrocephalus	+	-	-	+	-
3471	B	34	1	F T	Trial forceps failed	+	-	-	+	-
3592	B	29	1	F T	F.F.O.	+	-	-	+	-
3744	B	31	1	F T	C.P.	+	-	-	+	-
3770	A	34	4	F T	C.P.	+	-	-	+	-
3793	B	38	1	F T	Trial forceps failed	+	-	-	+	-
3868	B	18	1	F T	Malpresentation.	+	-	-	+	-
3973	B	38	6	F T	Impacted breach.	+	-	-	+	-
3987	B	26	1	F T	C.P.	+	-	-	+	-
4143	B	22	1	F T	C.P.	+	-	-	+	-
4252	B	38	6	F T	Hydrocephalus. Ruptured uterus	-	+	-	+	-
4205	B	34	5	F T	Hydrocephalus	+	-	-	+	-

6. *Oophorectomy, Ovariotomy, Salpingectomy,
and Salping-oophorectomy.*

The above operations were performed in 8 cases, as shown in the following table:—

Indication for Operation.	Mother.	
	Alive.	Died.
Ectopic pregnancy -	5	—
Ovarian cyst -	1	—
Bicornuate uterus -	1	—
Cystic ovary -	1	—
Totals - -	8	—

7. *Hysterectomy.*

Hysterectomy was performed in 5 cases, and the details of the cases are as follows:—

Reg. No.	Category.	Age.	Parity.	Stage of Pregnancy.	Indication for Operation.	Mother.		Child.		
						Alive.	Died.	Alive.	S.B.	Died.
158	A	29	3	F T	C.P. Ruptured uterus	+	—	—	+	—
188	B	39	3	F T	Ruptured uterus -	+	—	—	+	—
1044	A	35	7	—	Hydatid. mole -	+	—	—	—	—
2182	B	33	2	—	Conc. accid. hæm. -	+	—	—	—	—
2778	B	48	5	—	Ruptured uterus -	+	—	—	+	—
						5	—	—	3	—

8. *Hysterotomy.*

Abdominal hysterotomy was performed in 4 cases. Laparotomy and repair was performed in 1 case for ruptured uterus.

CONTRACTED PELVIS.

Three hundred and thirty-three cases of contracted pelvis were treated during the year 1930, and of these 208 were under ante-natal supervision, 103 were sent in as emergencies by outside doctors or midwives, or from the Outdoor Department, and 22 were admitted without having been under previous supervision.

Table—Cases of Contracted Pelvis which had been under Ante-Natal Supervision (Category A).

Mode of Delivery.	No. of Cases.	Mother.		Child.			Mortality per cent.	
		Alive.	Died.	Alive.	S.B.	Died.	M.	Ch.
Spontaneous - - -	65	65	—	61	2	2	—	3·0
Forceps - - -	32	31	1	23	8	1	3·1	25·0
Induction - - -	7	7	—	6 ¹	—	—	—	—
Cæsarean section - -	93	91	2	89	1	3	2·1	1·0
Craniotomy - - -	5	5	—	—	5	—	—	100·0
Version - - -	1	1	—	—	1	—	—	100·0
Undelivered - - -	2	2	—	—	—	—	—	—
Other obstetrical operations	3	3	—	—	3	—	—	100·0
Totals - - -	208	205	3	179	20	6	1·4	9·6

¹ 1 Case undelivered.

Cases sent in as Emergencies (Category B).

Mode of Delivery.	No. of Cases.	Mother.		Child.			Mortality per cent.	
		Alive.	Died.	Alive.	S.B.	Died.	M.	Ch.
Spontaneous - - -	25	25	—	21	4	—	—	16·0
Forceps - - -	39	36	3	27	12	1	7·6	30·7
Induction - - -	2	2	—	2	—	—	—	—
Cæsarean section - -	25	21	4	23	2	—	16·0	8·0
Craniotomy - - -	9	9	—	—	9	—	—	100·0
Version - - -	1	—	1	—	1	—	100·0	100·0
Undelivered - - -	2	2	—	—	—	—	—	—
Totals - - -	103	95	8	73	28	1	8·4	27·1

Note.—The percentage mortality for infants is calculated on the number of *stillbirths* amongst all children born, and does not take into account neonatal deaths.

Cases not under Ante-Natal Supervision—other than
Emergencies (Category C).

Mode of Delivery.	No. of Cases.	Mother.		Child.			Mortality per cent.	
		Alive.	Died.	Alive.	S.B.	Died.	M.	Ch.
Spontaneous - - -	7	7	—	5	1	1	—	14·2
Forceps - - -	4	4	—	4	—	—	—	—
Induction - - -	1	1	—	1	—	—	—	—
Cæsarean section - -	9	8	1	9	—	—	11·1	—
Craniotomy - - -	1	—	1	—	1	—	100·0	100·0
Totals - - -	22	20	2	19	2	1	9·0	9·0

Note.—The percentage mortality for infants is calculated on the number of *stillbirths* amongst all children born, and does not take into account neonatal deaths.

CONTRACTED PELVIS. 1. SPONTANEOUS DELIVERY.

A. Cases under Ante-natal Supervision.

Note.—The contractions G.C.P. denote Generally Contracted Pelvis : C.P., Contracted Pelvis; C.V., Conjugata Vera.

Reg. No.	Age.	Parity.	Maturity.	C.V.	Weight Child. ins. lbs.	Labour.		Mother.		Child.			Pyrexia.	Remarks.
						hrs.	min.	L.	D.	L.	D.	S.B.		
2	19	1	FT	3½	7¾	10	20	+	—	+	—	—	—	
10	26	3	FT	3½	7	10	15	+	—	+	—	—	—	
127	24	3	FT	4	7½	5	30	+	—	+	—	—	—	
166	25	3	FT	3+	6¼	9	7	+	—	+	—	—	—	
580	24	2	FT	3½	8	21	5	+	—	+	—	—	—	Successful trial labour
729	22	1	FT	3¾	8	8	30	+	—	+	—	—	—	
892	22	1	FT	3¼	6½	51	—	+	—	—	—	+	—	Rachitic flat pelvis. Hydrocephalic child
935	21	1	FT	3¾	7¼	28	20	+	—	+	—	—	—	
1086	24	1	FT	3½	6¾	13	10	+	—	+	—	—	—	
1096	23	1	FT	3¼	6½	32	—	+	—	+	—	—	—	Sl. rachitic-flat pelvis
1102	24	3	FT	3¼	8	4	35	+	—	+	—	—	—	Occ.-posterior
1150	23	1	FT	3¾	8½	8	20	+	—	+	—	—	—	Sl. G.C.P.
1175	—	—	FT	3½	7	17	—	+	—	+	—	—	—	Sl. C.P.
1186	24	4	FT	3¾	6½	12	45	+	—	+	—	—	—	Sl. G.C.P.
1197	22	2	FT	3¼+	6¼	8	40	+	—	+	—	—	—	
1221	24	1	FT	3¾	8	12	50	+	—	+	—	—	—	C. outlet, perineal tear
1286	26	1	FT	3¾	7	13	50	+	—	+	—	—	—	
1714	29	1	FT	—	7½	13	20	+	—	+	—	—	—	Sl. G.C.P. I.S. 9½" I.C. 10¼" E.C. 6¼"
1718	22	1	FT	3¼+	7	13	30	+	—	+	—	—	—	Rachitic G.C.P.
1728	17	1	FT	—	7¼	12	10	+	—	+	—	—	—	Sl. G.C.P.
1794	30	2	FT	3½+	6¾	18	20	+	—	+	—	—	—	Ant. parietal presentation
1809	25	1	FT	4	7	6	40	+	—	+	—	—	—	G.C.P. Contr. outlet. Complete tear
1831	19	2	FT	4	7½	19	30	+	—	+	—	—	—	
1833	27	4	FT	3½	6½	16	—	+	—	+	—	—	—	
1850	27	5	FT	3¾	9	14	12	+	—	+	—	—	—	
1871	20	1	FT	3½	6	18	50	+	—	+	—	—	—	
1878	22	1	FT	3¾	6¾	23	20	+	—	+	—	—	+	Sl. pyrexia three days
1882	22	1	FT	3½	6¾	8	30	+	—	+	—	—	—	
1887	25	1	FT	3¼	5¾	12	50	+	—	+	+	—	—	G.C.P. Child feeble

Reg. No.	Age.	Parity.	Maturity.	C.V.	Weight Child.	Labour.		Mother.		Child.			Pyrexia.	Remarks.
						hrs.	min.	L.	D.	L.	D.	S.B.		
1928	21	1	FT	3 $\frac{3}{4}$	8 $\frac{1}{2}$	14	10	+	—	+	—	—	—	Sl. pyrexia Outlet contracted G.C.P. Puerperal insanity on 5th day Transferred
1952	30	3	FT	3 $\frac{1}{2}$	6 $\frac{1}{2}$	12	35	+	—	+	—	—	—	
1953	20	1	FT	3 $\frac{3}{4}$	5 $\frac{1}{2}$	25	15	+	—	+	—	—	+	
2153	28	3	FT	—	8 $\frac{1}{2}$	24	—	+	—	+	—	—	—	
2161	22	1	FT	3 $\frac{3}{4}$	7 $\frac{1}{2}$	14	20	+	—	+	—	—	—	
2202	25	4	FT	3 $\frac{1}{4}$	7 $\frac{1}{2}$	12	25	+	—	+	—	—	—	Very little dis- proportion Patient small stature
2407	19	1	FT	3 $\frac{3}{4}$	7	6	40	+	—	+	—	—	—	
2675	25	3	FT	3 $\frac{3}{4}$	8	5	42	+	—	+	—	—	—	
2779	22	1	FT	3 $\frac{3}{4}$	6 $\frac{1}{2}$	12	50	+	—	+	—	—	—	
2803	25	3	FT	3 $\frac{3}{4}$	6 $\frac{1}{2}$	4	20	+	—	+	—	—	—	
2810	21	2	FT	3 $\frac{1}{2}$	6 $\frac{3}{4}$	6	30	+	—	+	—	—	—	Breech. Manual extraction.
2841	21	1	FT	3 $\frac{1}{2}$	7 $\frac{1}{2}$	12	20	+	—	+	—	—	—	
2960	21	2	FT	3 $\frac{3}{4}$	5 $\frac{1}{2}$	3	40	+	—	+	—	—	—	
3001	23	1	FT	4	8	11	25	+	—	+	—	—	—	Child feeble Child feeble
3046	32	3	FT	3 $\frac{3}{4}$	7	9	10	+	—	+	—	—	—	
3056	22	1	P	3 $\frac{1}{2}$	7 $\frac{3}{4}$	52	15	+	—	+	—	—	—	
3141	25	1	FT	3 $\frac{3}{4}$	6 $\frac{1}{2}$	10	20	+	—	+	—	—	—	Child feeble Child feeble Transverse at out- let, 2 $\frac{3}{4}$ "
3183	17	1	FT	3 $\frac{1}{2}$	6 $\frac{1}{2}$	29	20	+	—	+	—	—	—	
3214	27	6	FT	4	6 $\frac{1}{2}$	5	10	+	—	+	—	—	—	
3430	28	4	FT	3 $\frac{3}{4}$	9	10	20	+	—	+	—	—	—	Child feeble Child feeble W.R. + + Considerable œdema of vulva
3499	31	1	FT	3 $\frac{1}{2}$	7	9	35	+	—	+	—	—	—	
3584	24	2	FT	3 $\frac{1}{2}$	9 $\frac{1}{2}$	59	30	+	—	+	+	—	—	
3624	21	1	FT	4 $\frac{1}{4}$	5 $\frac{1}{2}$	12	20	+	—	+	—	—	—	Delivery after version
3773	34	6	FT	3 $\frac{1}{4}$	7 $\frac{1}{2}$	28	20	+	—	—	—	+	—	
3858	46	8	FT	3 $\frac{3}{4}$	7 $\frac{1}{4}$	9	5	+	—	+	—	—	—	
4074	34	7	FT	3 $\frac{1}{2}$	6	8	40	+	—	+	—	—	—	Rachitic flat pelvis
4159	28	2	FT	4—	8 $\frac{1}{4}$	5	30	+	—	+	—	—	—	
4174	23	1	FT	4—	7 $\frac{1}{4}$	17	10	+	—	+	—	—	—	
4191	23	1	FT	—	8	50	25	+	—	+	—	—	—	
4195	21	3	FT	3 $\frac{1}{4}$	6 $\frac{1}{2}$	12	40	+	—	+	—	—	—	
4291	29	2	FT	3 $\frac{3}{4}$	6 $\frac{3}{4}$	11	45	+	—	+	—	—	—	
4318	25	3	FT	3 $\frac{1}{2}$	8	17	50	+	—	+	—	—	—	
4329	19	1	FT	3 $\frac{3}{4}$	6	8	30	+	—	+	—	—	—	
4356	30	3	FT	3 $\frac{3}{4}$	7 $\frac{1}{4}$	3	15	+	—	+	—	—	—	
4428	23	1	FT	4	6	11	20	+	—	+	—	—	—	
4434	23	1	FT	—	—	—	—	+	—	+	—	—	—	

B. Cases sent in as Emergencies.

Reg. No.	Age.	Parity.	Maturity.	C.V.	Weight Child.	Labour.		Mother.		Child.			Pyrexia.	Remarks.
						hrs.	min.	L.	D.	L.	D.	S.B.		
707	21	1	FT	3 $\frac{3}{4}$	7	7	10	+	—	+	—	—	—	
708	28	1	FT	3	6 $\frac{1}{2}$	14	55	+	—	+	—	—	—	
1269	37	5	FT	3 $\frac{1}{4}$	7	4	35	+	—	+	—	—	—	
1411	32	1	FT	—	5 $\frac{3}{4}$	6	30	+	—	+	—	—	—	I.S. 8". I.C. 9 $\frac{1}{2}$ ". E.C. 6 $\frac{1}{2}$ "
1469	—	2	FT	4	8	26	30	+	—	+	—	—	—	
1646	21	2	FT	4—	7	7	35	+	—	+	—	—	+	Pyrexia only lasted 24 hours
1973	38	2	FT	4	6 $\frac{1}{4}$	12	35	+	—	+	—	—	—	
2038	24	2	FT	—	7	12	35	+	—	+	—	—	—	Feeble child. I.S. 9 $\frac{3}{4}$ ". I.C. 9 $\frac{3}{4}$ ". Rachitic pelvis
2249	23	2	FT	3 $\frac{3}{4}$	6	4	30	+	—	+	—	—	—	
2362	23	1	FT	4	7	11	10	+	—	—	—	+	—	Prol. cord
2535	22	1	FT	3 $\frac{1}{4}$	7 $\frac{1}{2}$	15	10	+	—	+	—	—	—	
2633	28	4	FT	3 $\frac{1}{2}$	5 $\frac{1}{2}$	46	—	+	—	—	—	+	—	Feeble child
3042	36	1	FT	—	—	—	—	+	—	+	—	—	—	
3115	26	2	FT	—	7	5	52	+	—	+	—	—	—	
3225	19	1	P	3 $\frac{1}{2}$	5 $\frac{3}{4}$	41	10	+	—	+	—	—	—	Premature child
3298	23	2	FT	3 $\frac{3}{4}$	7 $\frac{1}{2}$	3	35	+	—	+	—	—	—	
3309	25	3	FT	3 $\frac{3}{4}$	7	3	35	+	—	+	—	—	—	Transverse at out- let, 4" G.C.P.
3592	29	5	FT	—	7 $\frac{1}{4}$	12	10	+	—	+	—	—	—	Sl. disproportion
3747	24	1	FT	3 $\frac{3}{4}$	9	13	25	+	—	+	—	—	—	Large perineal tear
3755	20	1	FT	—	6 $\frac{1}{2}$	11	30	+	—	+	—	—	—	I.S. 9". I.C. 10". E.C. 7". Patient of small stature
3815	26	2	FT	3 $\frac{3}{4}$	7 $\frac{1}{2}$	14	10	+	—	+	—	—	—	I.S. 8 $\frac{3}{4}$ ". I.C. 9 $\frac{3}{4}$ "
3985	22	2	FT	—	7 $\frac{3}{4}$	21	20	+	—	+	—	—	—	I.S. 10". I.C. 10 $\frac{1}{2}$ " Sl. disproportion
3996	26	2	FT	—	8 $\frac{3}{4}$	25	5	+	—	—	—	+	—	Sl. disproportion
4151	24	2	FT	3 $\frac{1}{2}$	8	17	—	+	—	—	—	+	—	
4360	24	2	FT	3 $\frac{1}{4}$	7	8	—	+	—	+	—	—	—	

C. Not under Ante-natal Supervision.

793	22	1	FT	3 $\frac{1}{4}$	5 $\frac{1}{2}$	10	15	+	—	+	—	—	—	
1119	17	1	FT	3 $\frac{1}{4}$	6 $\frac{1}{4}$	5	15	+	—	+	—	—	—	
1205	20	1	FT	3 $\frac{3}{4}$	7 $\frac{1}{4}$	13	40	+	—	+	—	—	—	
1405	20	1	FT	4	6 $\frac{1}{2}$	21	15	+	—	—	—	—	—	
2975	20	1	FT	3 $\frac{1}{2}$	8	19	5	+	—	—	—	+	—	I.S. 9". I.C. 9 $\frac{1}{2}$ ". Trans. outlet, 4 $\frac{1}{4}$ " G.C.P. Gonorrhœa. Prol. cord
3662	19	1	FT	3 $\frac{3}{4}$	6 $\frac{1}{2}$	15	—	+	—	+	—	—	—	I.S. 8 $\frac{1}{2}$ ". I.C. 9 $\frac{1}{4}$ ". E.C. 6 $\frac{1}{2}$ "
3945	19	1	FT	3 $\frac{1}{4}$	7	21	—	+	—	+	+	—	—	Patient of small stature

CONTRACTED PELVIS. 2. FORCEPS DELIVERY
(NOT INDUCED).

A. Cases under Ante-natal Supervision.

Note.—F.F.O. denotes a case of Failed Forceps Outside of the Hospital practice.

Reg. No.	Age.	Parity.	Maturity.	C.V.	Weight Child. lbs.	Labour.		Mother.		Child.			Pyrexia.	Remarks.
						hrs.	min.	L.	D.	L.	D.	S.B.		
195	21	1	FT	4	7 $\frac{1}{4}$	15	0	+	—	+	—	—	—	Contracted outlet. Low forceps. Trans. at outlet, 3"
236	21	1	FT	—	8 $\frac{1}{2}$	31	—	+	—	+	—	—	—	Maternal distress. Sl. disproportion
687	30	4	FT	3 $\frac{3}{4}$	6 $\frac{1}{4}$	10	45	+	—	+	—	—	—	G.C.P.
747	43	1	FT	3 $\frac{1}{4}$	6	51	55	+	—	+	—	—	—	Foetal distress. Episiotomy
874	29	8	FT	3 $\frac{1}{2}$	8	46	20	+	—	+	—	—	—	Mid forceps Maternal distress
1428	18	1	FT	3 $\frac{1}{4}$	7 $\frac{3}{4}$	39	—	+	—	+	—	—	—	
1492	16	1	FT	3 $\frac{3}{4}$	6 $\frac{1}{2}$	12	20	+	—	+	—	—	—	Eclampsia com- plicating labour
1530	36	4	FT	4—	8 $\frac{1}{2}$	48	20	+	—	+	—	—	—	Maternal distress. High forceps
1570	18	1	FT	3 $\frac{3}{4}$	8	35	10	+	—	—	—	+	—	Foetal distress. Mid forceps
1666	28	1	FT	3 $\frac{3}{4}$	8	44	—	+	—	+	—	—	—	Maternal distress. G.C.P. Mid forceps
1764	34	1	FT	4—	4 $\frac{3}{4}$	32	30	+	—	+	—	—	+	High forceps. Mild pyrexia for 3 days
1815	18	1	FT	3 $\frac{1}{4}$	7 $\frac{3}{4}$	39	—	+	—	+	—	—	—	Maternal distress. Mid forceps
1905	23	1	FT	3 $\frac{3}{4}$	6 $\frac{3}{4}$	8	35	+	—	+	—	—	—	Sl. C.P. Spinal anæsthesia
2002	25	1	FT	3 $\frac{3}{4}$	8 $\frac{3}{4}$	16	0	+	—	+	—	—	—	Mid forceps. Sl. G.C.P.
2089	33	2	FT	4	8 $\frac{1}{2}$	6	50	+	—	+	—	—	—	Sl. contr. outlet. Mid forceps
2099	21	1	FT	3 $\frac{3}{4}$	6 $\frac{1}{4}$	17	20	+	—	—	—	+	—	Low forceps
2139	26	2	FT	3 $\frac{1}{2}$	6 $\frac{1}{2}$	12	55	+	—	+	—	—	—	Foetal distress. Mid forceps. Episiotomy
2265	25	1	FT	3 $\frac{3}{4}$	7 $\frac{1}{2}$	19	35	+	—	+	—	—	—	Maternal distress. Low forceps
2277	18	1	FT	3 $\frac{3}{4}$	8	31	—	+	—	—	—	+	—	Foetal distress. Mid forceps
2849	28	1	FT	4 $\frac{3}{4}$	7	10	40	+	—	+	—	—	—	Maternal distress. Low forceps. Large perineal tear
3113	21	1	FT	3 $\frac{3}{4}$	6 $\frac{3}{4}$	28	15	+	—	+	—	—	—	Foetal distress. Low forceps

Reg. No.	Age.	Parity.	Maturity.	C.V.	Weight Child. ins. lbs.	Labour.		Mother.		Child.			Pyrexia.	Remarks.
						hrs.	min.	L.	D.	L.	D.	S.B.		
3308	31	1	FT	3	—	19	5	+	—	+	+	—	—	Fœtal distress. Mid-forceps after trial labour
3365	23	1	FT	4	7 $\frac{3}{4}$	25	5	+	—	+	—	—	—	Fœtal distress. Low forceps
3409	23	1	FT	4	8 $\frac{3}{4}$	64	—	—	+	—	—	+	—	Low forceps. Obstet. shock
3412	22	1	FT	3 $\frac{1}{4}$	8	67	—	+	—	—	—	+	—	Maternal and fœtal distress. High forceps after trial labour
3467	22	1	FT	3 $\frac{1}{2}$	6	21	50	+	—	+	—	—	—	Mid forceps after trial labour. I.S. 8". I.C. 8 $\frac{1}{2}$ "
3507	24	1	FT	3 $\frac{3}{4}$	8 $\frac{1}{2}$	36	15	+	—	+	—	—	—	Fœtal distress. Low forceps
3539	22	1	FT	3 $\frac{1}{4}$	8	11	—	+	—	—	—	+	+	Trial labour. High Kielland forceps. Slight Naegele pelvis
3839	23	2	FT	3 $\frac{1}{2}$	7 $\frac{1}{2}$	16	40	+	—	—	—	+	—	Mid forceps
3852	—	—	FT	3 $\frac{3}{4}$	7 $\frac{1}{2}$	27	45	+	—	—	—	+	—	Mid forceps. Cord twisted round child's neck
4055	23	1	FT	4	6 $\frac{1}{2}$	24	—	+	—	+	—	—	—	Mid-forceps
4466	25	2	FT	3 $\frac{3}{4}$	8	25	—	+	—	+	—	—	—	High forceps

B. Cases sent in as Emergencies.

291	25	1	FT	3 $\frac{1}{2}$	7 $\frac{1}{2}$	6	25	—	+	+	—	—	+	Maternal distress. Died on 15th day. Pulmonary embolism
367	43	5	FT	3 $\frac{3}{4}$	9 $\frac{1}{2}$	72	30	+	—	—	—	+	—	Mid-forceps. G.C.P.
497	37	4	FT	—	—	27	—	+	—	+	—	—	—	C. outlet
625	27	1	FT	—	7 $\frac{3}{4}$	12	30	+	—	+	—	—	—	episiotomy
630	35	1	FT	—	8 $\frac{1}{4}$	62	—	+	—	+	—	—	—	Ant. parietal presentation
655	30	1	FT	3 $\frac{1}{2}$	7 $\frac{3}{4}$	15	—	+	—	—	—	+	—	F.F.O. High forceps
873	24	1	FT	3 $\frac{1}{4}$	7	49	15	+	—	+	—	—	+	Occ. post.
1012	24	1	FT	3 $\frac{1}{4}$	7 $\frac{3}{4}$	33	5	+	—	+	—	—	—	
1157	31	2	FT	—	6 $\frac{1}{2}$	28	10	+	—	—	—	+	—	G.C.P. Rachitic. Low forceps
1360	31	3	FT	—	6 $\frac{3}{4}$ & 8 $\frac{1}{2}$	15	55	—	+	—	—	++	+	F.F.O. Mid-forceps
1516	25	5	FT	3 $\frac{1}{2}$	7 $\frac{1}{2}$	66	45	+	—	—	—	+	—	G.C.P.
1531	21	1	FT	—	7 $\frac{1}{2}$	3	days	+	—	—	—	+	—	Marked disproportion
1609	24	1	FT	3 $\frac{1}{2}$	6	67	20	+	—	+	—	—	—	G.C.P. Mid-forceps

Reg. No.	Age.	Parity.	Maturity.	C.V.	Weight Child. ins. lbs.	Labour.		Mother.		Child.			Pyrexia.	Remarks.
						hrs.	min.	L.	D.	L.	D.	S.B.		
1722	24	1	FT	3 $\frac{1}{4}$	7 $\frac{1}{2}$	41	10	+	-	-	-	+	-	G.C.P. Mid-forceps
1842	32	1	FT	3 $\frac{3}{4}$	8 $\frac{1}{4}$	21	20	+	-	+	-	-	-	Low forceps.
														Contr. outlet
2208	34	2	FT	3 $\frac{3}{4}$	7 $\frac{3}{4}$	67	—	+	-	-	-	+	-	Low forceps
2017	28	1	FT	4	8 $\frac{1}{4}$	18	25	+	-	+	-	-	-	Post mature. C.
														outlet. 3 $\frac{1}{2}$ "
														transverse
2514	21	1	FT	3 $\frac{1}{2}$	10 $\frac{3}{4}$	4	days	-	+	+	-	-	+	Plac. manually
														removed. Adm.
														after 3 days'
														labour.
2648	25	1	FT	4	6	6	0	+	-	+	-	-	-	Low forceps
2748	23	1	FT	3 $\frac{1}{2}$	7 $\frac{3}{4}$	20	55	+	-	+	-	-	-	Mid-forceps.
														Foetal distress
2751	23	1	FT	3 $\frac{3}{4}$	7 $\frac{1}{4}$	21	35	+	-	+	-	-	-	Foetal distress.
														Low forceps
2864	24	1	FT	3 $\frac{3}{4}$	7	17	5	+	-	+	-	-	+	Mid-forceps
														P.P.H.
2870	32	3	FT	3 $\frac{1}{2}$	9	11	10	+	-	+	-	-	-	Low forceps
2958	40	1	FT	3 $\frac{3}{4}$	7 $\frac{1}{2}$	39	0	+	-	+	-	-	-	Low forceps
2988	29	5	FT	3 $\frac{3}{4}$	7	50	45	+	-	+	-	-	-	Mid-forceps.
														Maternal distress
2990	24	1	FT	3 $\frac{1}{2}$	7 $\frac{1}{4}$	28	35	+	-	-	-	+	-	Low forceps
														Foetal distress
3049	35	1	FT	3 $\frac{3}{4}$	5 $\frac{1}{2}$	11	30	+	-	+	-	-	-	Low forceps
3088	18	1	FT	3 $\frac{1}{2}$	7 $\frac{1}{2}$	19	40	+	-	+	-	-	+	Mid-forceps.
														Maternal distress
3375	32	1	FT	4	6 $\frac{1}{4}$	26	—	+	-	-	-	+	-	Mid-forceps
3380	28	5	FT	3 $\frac{3}{4}$	10 $\frac{1}{2}$	36	0	+	-	+	-	-	+	Rachitic flat pel-
														vis. High forceps
3722	22	1	P	3 $\frac{1}{4}$	5 $\frac{1}{4}$	22	—	+	-	+	+	-	-	High forceps
														Feeble child
3931	34	1	FT	—	7 $\frac{1}{2}$	36	—	+	-	+	-	-	-	Low forceps.
														Feeble child
														Sl. disproportion
4028	26	1	FT	3 $\frac{3}{4}$ +	6 $\frac{3}{4}$	11	50	+	-	+	-	-	-	Low forceps.
														F.F.O.
4036	27	1	FT	3 $\frac{1}{2}$	5 $\frac{1}{4}$	36	10	+	-	+	-	-	-	Mid-forceps.
														Maternal distress
4077	29	4	FT	—	8 $\frac{3}{4}$	36	15	+	-	+	-	-	-	Sl. disproportion
4142	30	5	FT	3 $\frac{3}{4}$	9 $\frac{1}{2}$	50	—	+	-	+	-	-	+	High forceps
4258	30	3	FT	4	8 $\frac{3}{4}$	18	20	+	-	+	-	-	-	Maternal distress.
														Mid-forceps
4320	33	3	FT	—	8 $\frac{1}{4}$	8	—	+	-	+	-	-	-	Low forceps
4330	22	1	FT	3 $\frac{1}{4}$ +	6 $\frac{3}{4}$	23	—	+	-	+	-	-	-	Mid-forceps
C. Cases not under Ante-natal Supervision.														
506	20	1	FT	3 $\frac{1}{2}$	7	6	25	+	-	+	-	-	-	Mid-forceps.
														G.C.P.
1413	23	2	FT	3 $\frac{1}{2}$	6 $\frac{3}{4}$	19	20	+	-	+	-	-	-	Low forceps
														Foetal distress
3487	22	1	FT	4	8	7	50	+	-	+	-	-	-	Low forceps
4140	38	1	FT	3 $\frac{1}{2}$	6 $\frac{1}{2}$	48	15	+	-	+	-	-	-	Low forceps
														Foetal distress

CONTRACTED PELVIS. 3. INDUCTION OF LABOUR.

A.—Cases under Ante-natal Supervision.

Reg. No.	Age.	Parity.	Maturity.	C.V.	Weight Child.	Labour.		Mother.		Child.			Pyrexia.	Remarks.
						hrs.	min.	L.	D.	L.	D.	S.B.		
937	30	3	FT	3¾	7¾	6	45	+	—	+	—	—	—	Epilepsy Med. induction
938	25	2	—	3¾	—	—	—	+	—	—	—	—	—	Med. ind. unsuccessful
1375	21	1	FT	—	8¼	15	45	+	—	+	—	—	—	Med. ind. at term. Sl. disproportion
1894	23	1	FT	3¼	5¾	3	25	+	—	+	—	—	+	Med. ind. at 37th week
1958	26	1	FT	—	9	5	15	+	—	+	—	—	—	Med. ind. at 38th week. Contr. outlet
2547	30	4	FT	3¾	6½	53	—	+	—	+	—	—	—	Med. ind. at term
4202	28	3	FT	—	8¾	6	15	+	—	+	—	—	—	Sl. C.P. Med Ind. at term

B.—Cases sent in as Emergencies.

2642	32	2	FT	3¼	9	4	5	+	—	+	—	—	—	Med. Ind. at term
4253	29	1	P	—	6¾	7	15	+	—	+	—	—	—	Sl. C.P. surgical Ind. (Bougies) at 38th week

C.—Cases not under Ante-natal Supervision.

2058	18	1	FT	3¼	6	6	40	+	—	+	—	—	—	Med. Ind. at term
------	----	---	----	----	---	---	----	---	---	---	---	---	---	-------------------

CONTRACTED PELVIS. 4. CÆSAREAN SECTION.

A. Cases under Ante-natal Supervision.

Note.—L.U.S. denotes Lower Uterine Segment operation. The number in brackets after "Repeat section" denotes the number of sections performed.

207	21	1	FT	—	6¼	—	—	+	—	+	—	—	—	W.R. + +
208	25	1	FT	3	7¾	—	—	+	—	+	—	—	+	Urinary infection
267	37	4	FT	3	7½	5	—	+	—	+	—	—	—	Repeat section (2). Sterilized
359	21	1	FT	3	7	6	—	+	—	+	—	—	—	L.U.S.
514	25	2	FT	3¼	9	—	—	+	—	+	—	—	—	
640	24	1	FT	2¾	6¾	—	—	+	—	+	—	—	—	Sterilized
785	27	3	FT	3	9	—	—	—	+	—	+	—	—	Repeat section (2). Sterilized. Spinal anæsthesia
811	26	1	FT	—	8	10	—	+	—	+	—	—	+	L.U.S. Trans. at outlet, 2½"

Reg. No.	Age.	Parity.	Maturity.	C.V.	Weight Child. ins. lbs.	Labour.		Mother.		Child.			Pyrexia.	Remarks.
						hrs.	min.	L.	D.	L.	D.	S.B.		
882	20	1	FT	3	7½	—	—	+	—	+	—	—	—	Acute intestinal obstruction two months later
930	37	3	FT	3¾	8½	—	—	+	—	+	—	—	—	Bad obstet. hist.
977	26	3	FT	3¼	6¼	—	—	+	—	+	—	—	—	Bad obstet. hist.
1005	23	1	FT	3⅛	6½	12	—	+	—	+	—	—	+	L.U.S. Mild pyrexia
1129	27	2	FT	—	7	—	—	+	—	+	—	—	—	Spinal anæsthesia. Repeat section (2). Sterilized
1141	21	1	—	2¾	6	3	—	+	—	+	—	—	—	Sterilized
1146	26	2	FT	—	10½	1	45	+	—	+	—	—	—	Repeat section (2) Sterilized
1170	31	3	FT	3¼	7	—	—	+	—	+	—	—	—	Repeat section (3). Sterilized
1172	29	1	FT	2¼	8	—	—	+	—	+	—	—	—	
1174	24	3	FT	2¾	5¾	8	—	+	—	+	—	—	—	Repeat section (2) Spinal anæsthesia
1199	30	1	FT	3½	6¼	—	—	+	—	+	—	—	—	
1232	25	2	FT	4	8¼	—	—	+	—	+	—	—	—	Sterilized. Repeat section (2). Rachitic, flat
1243	27	2	FT	3¾	6½	12	—	+	—	+	—	—	+	Marked disproportion. L.U.S.
1252	40	2	FT	3	8	—	—	+	—	+	—	—	+	Breast abscess Repeat section (2). Sterilization
1327	35	4	FT	3	6½	—	—	+	—	+	—	—	—	Sterilized
1367	29	1	FT	—	7	—	—	+	—	+	—	—	—	G.C.P.
1383	26	2	FT	3¼	7¼	—	—	+	—	+	—	—	—	Repeat section. Spinal anæsthesia
1384	33	2	FT	—	8¼	—	—	+	—	+	—	—	—	Marked disproportion
1453	38	1	FT	—	7¼	7	—	+	—	+	—	—	—	L.U.S. Marked disproportion.
1557	31	6	FT	3¼	7	—	—	+	—	+	—	—	—	Sterilized
1565	28	1	FT	3¼	7¼	2	—	+	—	+	—	—	—	Spinal anæsthesia
1566	25	3	FT	—	5¾	—	—	+	—	+	—	—	—	Spinal anæsthesia. Repeat section (3). Sterilized
1581	30	1	FT	3¼	6¾	7	—	+	—	+	—	—	—	L.U.S.
1628	25	1	FT	3¼	10½	12	—	+	—	+	—	—	—	L.U.S.
1641	35	4	FT	3	8	—	—	+	—	+	—	—	—	Sterilized
1653	29	4	P	3¾	8	—	—	+	—	+	—	—	—	Repeat section (3) Sterilized.
1655	23	3	FT	3¼	8½	8	—	+	—	+	—	—	—	Spinal anæsthetic L.U.S. Spinal anæsthesia

Reg. No.	Age.	Parity.	Maturity.	C.V. ins.	Weight Child. lbs.	Labour.		Mother.		Child.			Pyrexia.	Remarks.
						hrs.	min.	L.	D.	L.	D.	S.B.		
1832	30	4	FT	3 $\frac{1}{4}$	5 $\frac{1}{4}$ & 6	—	—	+	—	+	—	—	—	Marked rachitic pelvis
1897	38	1	FT	—	7	—	—	+	—	+	—	—	—	Repeat section (2) G.C.P. and L.U.S.
1924	29	2	FT	3	7	—	—	+	—	+	—	—	—	Sterilized Repeat section (2)
1959	35	3	FT	2	8 $\frac{3}{4}$	—	—	+	—	+	—	+	—	Repeat section (3) Sterilized. Achondroplasia
2005	22	1	FT	3	7 $\frac{1}{4}$	—	—	+	—	+	—	—	—	Repeat section (2) Spinal anæ- sthetic. Rachitic pelvis. Marked disproportion
2039	24	2	FT	—	9	—	—	+	—	+	—	—	—	
2068	36	2	FT	—	11	—	—	+	—	+	—	—	—	Spinal anæsthesia Marked dispro- portion
2151	18	1	FT	—	7 $\frac{1}{4}$	—	—	+	—	+	—	—	—	Marked disproportion
2163	27	4	FT	2 $\frac{3}{4}$	9 $\frac{1}{4}$	—	—	+	—	+	—	—	—	Sterilized. Bad obstet. hist.
2170	24	3	FT	3 $\frac{1}{2}$	8	—	—	+	—	+	—	—	—	Spinal anæ- sthetic Bad obstet. hist.
2191	22	1	FT	2 $\frac{1}{4}$	7	4	—	+	—	+	—	—	—	Marked rachitic pelvis
2239	31	4	FT	—	8 $\frac{1}{4}$	—	—	+	—	+	—	—	—	Spinal anæsthesia. Repeat section (2). Sterilized
2316	27	5	FT	3 $\frac{1}{4}$	8	10	—	+	—	+	—	—	+	Sterilized. Bad obstet. hist.
2327	22	1	FT	3 $\frac{3}{4}$	5 $\frac{3}{4}$	3 $\frac{1}{2}$	—	+	—	+	—	—	—	L.U.S.
2367	30	4	FT	3 $\frac{1}{4}$	8	48	—	+	—	+	—	—	—	Sterilized
2419	30	1	FT	3 $\frac{1}{2}$	5 $\frac{3}{4}$	12	—	+	—	+	—	—	+	G.C.P. and L.U.S.
2729	33	2	FT	—	6 $\frac{1}{2}$	4	—	+	—	+	—	—	—	Repeat section (2)
2777	24	1	FT	3	8 $\frac{1}{2}$	—	—	+	—	+	—	—	+	Rachitic
2788	24	1	FT	3	6 $\frac{3}{4}$	12	—	+	—	+	—	—	—	L.U.S. Spinal anæsthesia
2858	24	1	P	3 $\frac{1}{4}$	6	7	—	+	—	+	—	—	—	Spinal anæs- thetic. L.U.S.
2999	39	2	FT	3	7 $\frac{3}{4}$	—	—	+	—	+	—	—	—	Repeat section (2) Sterilized
3048	24	2	FT	2 $\frac{1}{2}$	9 $\frac{3}{4}$	2	—	+	—	+	—	—	—	Sterilized. Repeat section (2)
3052	25	3	FT	3 $\frac{1}{2}$	6	4	—	+	—	+	—	—	—	L.U.S.
3204	30	5	P	—	5 $\frac{3}{4}$	—	—	+	—	+	—	—	—	Repeat section (3) Sterilized

Reg. No.	Age.	Parity.	Maturity.	C.V. ins.	Weight Child. lbs.	Labour.		Mother.		Child.			Pyrexia.	Remarks.
						hrs.	min.	L.	D.	L.	D.	S.B.		
3268	30	2	FT	3	8	—	—	+	—	+	—	—	—	Repeat section (2) W.R. + + Sterilized
3280	25	3	FT	3 $\frac{1}{4}$	5 $\frac{3}{4}$	—	—	—	+	+	—	—	—	Repeat section (2) Anæsthetic death
3290	27	1	FT	3 $\frac{1}{4}$	6 $\frac{1}{2}$	24	—	+	—	+	—	—	—	L.U.S.
3353	32	4	FT	3 $\frac{1}{2}$	8 $\frac{1}{2}$	4	—	+	—	+	—	—	—	Repeat section (2). Sterilized
3408	32	4	FT	3 $\frac{1}{4}$	8 $\frac{1}{2}$	—	—	+	—	+	—	—	—	G.C.P. Bad obstet. hist.
3421	—	6	FT	3 $\frac{1}{2}$	9	—	—	+	—	+	—	—	—	Repeat section (3) Sterilized. Spinal anæsthetic
3426	31	5	FT	3 $\frac{1}{4}$	6 $\frac{1}{2}$	6	—	+	—	+	—	—	—	Repeat section (2) L.U.S. Spinal anæsthetic. Sterilized.
3451	24	1	FT	3	6 $\frac{1}{2}$	72	—	+	—	+	+	—	—	Section after trial labour
3481	18	1	FT	3	8	9	—	+	—	+	—	—	—	L.U.S.
3483	20	1	FT	3 $\frac{1}{2}$	6 $\frac{1}{2}$	8	20	+	—	+	—	—	—	L.U.S.
3514	21	1	FT	3 $\frac{1}{4}$	6 $\frac{1}{4}$	27	—	+	—	+	—	—	+	
3580	21	1	FT	2 $\frac{1}{2}$	6	—	—	+	—	+	+	—	—	L.U.S.
3598	24	1	FT	3	7 $\frac{1}{2}$	6	—	+	—	+	—	—	—	
3613	25	1	FT	3 $\frac{3}{4}$	7 $\frac{1}{2}$	10	—	+	—	+	—	—	+	Spinal anæsthetic Med. induct. L.U.S.
3680	28	2	FT	2 $\frac{3}{4}$	7 $\frac{1}{2}$	3	—	+	—	+	—	—	—	Marked rachitic deformities. Sterilized
3681	22	3	FT	2 $\frac{3}{4}$	7	4	—	+	—	+	—	—	—	Repeat section (2) Sterilized
3682	28	2	FT	—	5 $\frac{3}{4}$	—	—	+	—	+	—	—	—	Sterilized. Marked dis- proportion. Rachitic defor- mities
3690	27	2	FT	3 $\frac{1}{4}$	7 $\frac{1}{4}$	—	—	+	—	+	—	—	—	
3699	19	1	FT	3 $\frac{1}{4}$	8 $\frac{1}{2}$	2	30	+	—	+	—	—	—	
3841	27	1	FT	—	7 $\frac{1}{2}$	48	—	+	—	—	—	+	—	Marked dispropor- tion. Rachitic flat pelvis
3883	24	2	FT	3 $\frac{1}{4}$	7 $\frac{1}{4}$	3	15	+	—	+	—	—	+	Sterilized. Spinal anæsthetic
3896	29	2	FT	—	7 $\frac{3}{4}$	12	15	+	—	+	—	—	—	Marked disproportion L.U.S.
3903	25	2	FT	3 $\frac{1}{2}$	7	5	—	+	—	+	—	—	+	
3993	26	1	FT	3 $\frac{1}{4}$	6 $\frac{1}{2}$	3	10	+	—	+	—	—	+	L.U.S.
4061	21	1	FT	3	7	—	—	+	—	+	—	—	—	Spinal anæsthetic

Reg. No.	Age.	Parity.	Maturity.	C V.	Weight Child. ins. lbs.	Labour.		Mother.		Child.			Pyrexia.	Remarks.
						hrs.	mins.	L.	D.	L.	D.	S.B.		
4081	29	2	P	3	4½	—	—	+	—	+	—	—	—	Pains for 1 hour 35 mins.
4136	24	2	FT	—	6½	4	20	+	—	+	—	—	—	Repeat section (2) Sterilized
4167	25	2	FT	—	6½	—	—	+	—	+	—	—	—	
4189	23	1	FT	3	7½	3	—	+	—	+	—	—	—	Spinal anæsthetic. L.U.S.
4201	30	6	FT	2½	6½	—	—	+	—	+	—	—	—	Repeat section (3) Sterilized
4348	24	1	FT	3½	7	12	—	+	—	+	—	—	—	L.U.S.
4389	18	1	FT	3	8	—	—	+	—	+	—	—	—	Spinal anæsthetic
4395	30	5	FT	—	7	—	—	+	—	+	—	—	—	Med. ind. Sterilized. Bad obstet. hist.
4446	25	2	FT	3¼	8½	—	—	+	—	+	—	—	—	Repeat section (2) Sterilized

B. Cases sent in as Emergencies.

202	24	2	FT	3½	8	4	50	+	—	+	—	—	—	G.C.P.
249	25	1	FT	2¼	8¼	—	—	+	—	+	—	—	—	Spinal bifida
542	24	1	FT	2¾	7¾	36	—	—	+	+	—	—	—	Patient was per- suaded with diffi- culty to come into hospital.
669	30	1	FT	3	6¼	48	—	+	—	—	—	+	—	Multiple vaginal examinations outside. No foetal heart on admission
718	22	1	FT	3¼	6	8	35	+	—	+	—	—	—	L.U.S. Spinal anæsthesia
881	37	1	FT	2¾	7½	9	20	+	—	+	—	—	+	Good recovery
1268	35	2	FT	3½	7¾	12	—	+	—	+	—	—	—	L.U.S. Spinal anæ. G.C.P.
1363	33	1	FT	—	8	6	—	+	—	+	—	—	—	L.U.S.
1370	27	6	FT	—	7¼	7	55	+	—	+	—	—	—	Sterilized. Bad obstet. hist.
1385	20	1	FT	3	6½	5	40	+	—	+	—	—	+	Spinal anæsthetic
1606	30	1	FT	3	8	9	—	+	—	+	—	—	+	L.U.S.
1665	25	2	FT	3	7½	5	15	+	—	+	—	—	+	L.U.S.
1899	27	1	FT	3	6¾	—	—	+	—	+	—	—	—	
2049	25	2	FT	—	6¼	2	35	+	—	+	—	—	—	
2350	25	3	FT	3¾	6	36	—	+	—	+	—	—	+	Repeat section (3) Sterilized
2482	26	1	FT	3¼	8	—	—	+	—	+	—	—	+	
2874	31	1	FT	—	9¼	2	30	+	—	+	—	—	—	Marked dispro- portion. Myo- mectomy

Reg. No.	Age.	Parity.	Maturity.	C.V.	Weight Child.	Labour.		Mother.		Child.			Pyrexia	Remarks.
						hrs.	mins.	L.	D.	L.	D.	S.B.		
3101	25	1	FT	3	8	7	45	—	+	+	—	—	+	Peritonitis.
3292	25	1	FT	3	6	12	50	—	+	+	—	—	—	Left basal
3805	26	4	FT	3	8	23	—	—	+	—	—	+	+	P.M. Pneumonia.
														L.U.S. (F.F.O.)
														Peritonitis
3938	20	1	FT	3 $\frac{1}{4}$	6 $\frac{1}{2}$	—	—	+	—	+	—	—	—	
3981	26	4	FT	3 $\frac{1}{4}$	7 $\frac{3}{4}$	—	—	+	—	+	—	—	—	Spinal anæ-
														thetic and bad
														obstet. hist.
4070	23	1	FT	3 $\frac{1}{2}$	7 $\frac{1}{2}$	25	—	+	—	+	—	—	—	L.U.S.
4314	37	10	FT	3 $\frac{1}{4}$	8	36	—	+	—	+	—	—	—	Spinal
														anæsthetic
4467	18	1	FT	—	6 $\frac{3}{4}$	7	—	+	—	+	—	—	—	Spinal
														anæsthetic
														L.U.S.

C. Cases not under Ante-natal Supervision.

537	23	1	FT	3—	8	8	—	+	—	+	—	—	—	
950	23	3	FT	—	6 $\frac{3}{4}$	4	25	+	—	+	—	—	—	Repeat section (3)
														Sterilized.
														Spinal
														anæsthetic
2606	30	3	FT	3 $\frac{1}{2}$	8 $\frac{1}{2}$	12	—	+	—	+	—	—	—	Repeat section (3)
														Sterilized.
2931	29	3	P	2 $\frac{3}{4}$	4 $\frac{3}{4}$	4	—	—	+	+	—	—	—	Pulmonary
														embolism
3171	24	2	FT	3	6 $\frac{3}{4}$	—	—	+	—	+	—	—	+	Repeat section (2)
														Sterilized
3487	22	1	FT	3 $\frac{1}{4}$	6 $\frac{1}{2}$	2	—	+	—	+	—	—	—	
3647	22	1	FT	3	6 $\frac{1}{4}$	2	—	+	—	+	—	—	—	
3795	40	2	FT	3 $\frac{1}{2}$	7	—	—	+	—	+	—	—	—	
4170	27	2	FT	3 $\frac{1}{4}$	8	6	—	+	—	+	—	—	—	

CONTRACTED PELVIS. 5. CRANIOTOMY.

A. Cases under Ante-natal Supervision.

25	39	4	FT	—	—	—	—	+	—	—	—	+	—	
69	23	1	FT	3	6	58	55	+	—	—	—	+	—	
191	40	5	FT	3 $\frac{1}{2}$	6 $\frac{1}{2}$	10	25	+	—	—	—	+	—	F.F.
546	32	4	FT	2 $\frac{3}{4}$	5 $\frac{1}{2}$	12	40	+	—	—	—	+	—	Posterior parietal
														Prol. cord
3770	34	4	FT	3 $\frac{1}{4}$	7 $\frac{1}{2}$	31	—	+	—	—	—	+	—	F.F. Bad obstet.
														hist.

B. Cases sent in as Emergencies.

Reg. No.	Age.	Parity.	Maturity.	C.V.	Weight Child. ins. lbs.	Labour.		Mother.		Child.			Pyrexia.	Remarks.
						hrs.	mins.	L.	D.	L.	D.	S.B.		
555	27	1	FT	3 $\frac{3}{4}$	7 $\frac{1}{2}$	9+	—	+	—	—	—	+	—	F.F.O.
1305	35	1	FT	3 $\frac{1}{2}$	9	36	—	+	—	—	—	+	—	Delay in 1st stage
1636	23	1	FT	3 $\frac{1}{4}$	6 $\frac{1}{4}$	42	—	+	—	—	—	+	—	F. Forceps
2645	31	2	FT	3 $\frac{1}{2}$	8	36	—	+	—	—	—	+	—	Severe cardiac disease
3744	31	1	FT	3 $\frac{1}{4}$	8 $\frac{1}{2}$	44	—	+	—	—	—	+	+	Failed forceps
3918	25	1	FT	3 $\frac{3}{4}$	—	20	30	+	—	—	—	+	—	Hydrocephalic head
3987	26	1	FT	3	6 $\frac{3}{4}$	14	—	+	—	—	—	+	—	Breech. Head perforated
4143	22	1	FT	3 $\frac{1}{2}$	7	11	45	+	—	—	—	+	—	Prol. cord. Head perforated
4269	30	2	FT	3 $\frac{1}{4}$	6 $\frac{1}{2}$	36	—	+	—	—	—	+	—	Previous outside interference

C. Cases not under Ante-natal Supervision.

3797	26	4	FT	3 $\frac{1}{4}$	7	12	—	—	+	—	—	+	+	Ruptured uterus
------	----	---	----	-----------------	---	----	---	---	---	---	---	---	---	-----------------

CONTRACTED PELVIS. 6. VERSION.

A. Cases under Ante-natal Supervision.

4174	23	1	FT	3 $\frac{3}{4}$	7 $\frac{1}{4}$	17	10	+	—	+	—	—	—	Ext. version from breech before labour commenced
------	----	---	----	-----------------	-----------------	----	----	---	---	---	---	---	---	--

B. Cases sent in as Emergencies.

3753	34	1	FT	—	9 $\frac{1}{4}$	48	—	—	+	—	—	+	—	Marked disproportion. F.F. Int. podalic version
------	----	---	----	---	-----------------	----	---	---	---	---	---	---	---	---

C. Cases not under Ante-natal Supervision.

Nil.

CONTRACTED PELVIS. 7. NOT DELIVERED.

A. Cases under Ante-natal Supervision.

Reg. No.	Age.	Parity.	Maturity.	C.V.	Weight Child.	Labour.		Mother.		Child.			Pyrexia.	Remarks.
						hrs.	mins.	L.	D.	L.	D.	S.B.		
255	44	12	P	3 $\frac{1}{4}$	—	—	—	+	—	—	—	—	—	No marked disproportion Adm. with pyelitis
2088	17	2	P	3 $\frac{1}{4}$	—	—	—	+	—	—	—	—	—	

B. Cases sent in as Emergencies.

773	31	2	—	—	—	—	—	+	—	—	—	—	—	G.C.P. Non-pregnant Incom. abortion
1992	29	1	3	2 $\frac{3}{4}$	—	—	—	+	—	—	—	—	—	

C. Cases not under Ante-natal Supervision.

Nil.

CONTRACTED PELVIS. 8. OTHER OBSTETRICAL OPERATIONS.

A. Cases under Ante-natal Supervision.

157	29	3	FT	3	7	—	—	+	—	—	—	+	—	Adm. with rupt. uterus. Sub-total hysterectomy Prim. breech (impacted), manual extraction Manual extraction of breech. Puer. sepsis. Transferred to fever hospital
183	25	1	FT	3 $\frac{1}{2}$	7	9	10	+	—	—	—	+	—	
438	30	3	FT	3 $\frac{1}{4}$	7 $\frac{1}{2}$	3	25	+	—	—	—	+	+	

B. Cases sent in as Emergencies.

Nil.

C. Cases not under Ante-natal Supervision.

Nil.

ANTE-PARTUM HÆMORRHAGE.

There were 206 cases of ante-partum hæmorrhage admitted during the year, and of these 94 were cases of placenta prævia and 112 were cases of accidental hæmorrhage.

1. PLACENTA PRÆVIA.

A. Cases under Ante-natal Supervision (Total, 18).

Method of Treatment.	Type.		Mother.		Child.			Mortality. per cent.	
	Cent.	Lat.	A.	D.	A.	S.B.	D.	Mother.	Child.
Spontaneous delivery, - -	—	7	7	—	6	1	—	—	14·2
Cæsarean section, - - -	1	3	4	—	4	—	—	—	—
Version, - - - -	1	4	5	—	2	3	—	—	60·0
Manual delivery, - - -	—	2	2	—	—	1	1	—	50·0
Totals, - - -	2	16	18	—	12	5	1	—	27·7

B. Cases sent in as Emergencies (Total, 64).

Spontaneous delivery, - -	1	10	11	—	8	3	—	—	—
Cæsarean section, - - -	16	4	19	1	13	2	5	5·0	—
Version, - - - -	14	11	23	2	4	20	1	8·0	—
Forceps, - - - -	—	1	1	—	—	1	—	—	—
Packing - - - -	2	4	4	2 ¹	3 ²	2	—	33·3	—
Manual delivery - - -	—	1	1	—	—	1	—	—	—
Total, - - -	33	31	59	5	28	29	6	7·8	—

¹ Two mothers died undelivered. ² One case of twins.

C. Cases not under Ante-natal Supervision (Total, 12).

Method of Treatment.	Type.		Mother.		Child.			Mortality per cent.	
	Cent.	Lat.	A.	D.	A.	S.B.	D.	Mother	Child.
Spontaneous delivery, - - -	—	5	5	—	3	2	—	—	40·0
Cæsarean section, - - -	3	2	4	1	1	—	5 ¹	20·0	—
Version, - - - -	—	1	—	1	—	1	—	100·0	100·0
Packing, - - - -	—	1	1	—	—	1	—	—	100·0
Totals, - - -	3	9	10	2	4	4	5	16·6	33·3

¹ One case of twins.

All Groups (A, B, and C) (Total, 94).

Spontaneous delivery, - - -	1	22	23	—	17	6	—	—	26·0
Cæsarean section, - - -	20	9	27	2	18	2	10	6·8	6·8
Version, - - - -	15	16	28	3	6	24	1	9·6	77·4
Forceps, - - - -	—	1	1	—	—	1	—	—	100·0
Packing, - - - -	2	5	5	2	3	3	—	28·5	50·0
Manual delivery - - -	—	3	3	—	—	2	1	—	66·6
Totals, - - -	38	56	87	7	44	38	12	7·4	40·4

2. ACCIDENTAL HÆMORRHAGE.

A. Cases under Ante-natal Supervision (Total, 16).

Method of Treatment.	Type.			Mother.		Child.			Mortality.	
	Con- cealed.	Appar- ent.	Com- bined.	A.	D.	A.	S.B.	D.	Mother.	Child.
Spontaneous delivery, -	—	10	2	12	—	7	5	—	—	41·6
Forceps, - - -	—	2	—	2	—	—	1	1	—	50·0
Undelivered, - -	1	1	—	1	1	—	—	—	50·0	—
Totals, - -	1	13	2	15	1	7	6	1	5·8	75·0

B. Cases sent in as Emergencies (Total, 81).

Spontaneous delivery, -	11	29	20	58	2	25 ¹	35	4	3·2	57·3
Cæsarean section, -	—	1	1	2	—	—	2	—	—	100·0
Forceps, - - -	1	1	1	2	1	—	3	—	33·3	100·0
Version, - - -	—	1	2	3	—	2	1	—	—	33·3
Packing, - - -	—	3	—	3	—	2	1	—	—	33·3
Hysterectomy, - -	1	—	—	1	—	—	1	—	—	100·0
Manual delivery, - -	1	—	1	2	—	—	2	—	—	100·0
Undelivered, - -	3	1	3	1	6	—	—	—	—	—
Totals, - -	17	36	28	72	9	29	45	4	10·9	56·0

¹ Twins (4 times).

C. Cases not under Ante-natal Supervision (Total, 15).

Method of Treatment.	Type.			Mother.		Child.			Mortality.	
	Con- cealed.	Appar- ent.	Com- bined.	A.	D.	A.	S.B.	D.	Mother.	Child.
Spontaneous delivery, -	2	8	3	12	1	6	4	3	7·6	30·7
Undelivered, - - -	—	2	—	2	—	—	—	—	—	—
Totals, - - -	2	10	3	14	1	6	4	3	7·1	3·7

All Groups (A, B, and C) (Total, 112).

Spontaneous delivery, -	13	47	25	82	3	38 ¹	44	7	3·5	48·8
Cæsarean section, -	—	1	1	2	—	—	2	—	—	100·0
Forceps, - - -	1	3	1	4	1	—	4	1	—	80·0
Version, - - -	—	1	2	3	—	2	1	—	—	33·3
Packing, - - -	—	3	—	3	—	2	1	—	—	33·3
Manual delivery, - -	1	—	1	2	—	—	2	—	—	100·0
Hysterectomy, - -	1	—	—	1	—	—	1	—	—	100·0
Undelivered, - -	4	4	3	4	7	—	—	—	58·3	—
Totals, - - -	20	59	33	101	11	42	55	8	9·8	49·1

¹ Twins (4 times).

Details of the cases of ante-partum hæmorrhage are shown in the following tables:—

PLACENTA PRÆVIA.

A. Cases under Ante-natal Supervision.

Reg. No.	Age.	Parity.	Maturity.	Condition on Admission.	Type.		Treatment.	Result.		Remarks.
					Cnt.	Lat.		M.	Ch.	
381	34	7	FT	Fairly good	—	+	Packing, spontaneous delivery	A	A	
633	30	10	P	Fair, intermittent bleeding	—	+	Spontaneous delivery	A	A	High lat. type
1006	32	3	FT	Poor, severe bleeding	+	—	Podalic version	A	SB	Puer. septicæmia Transferred to Belvedere Hospital
1125	31	5	FT	Good	—	+	Podalic version	A	A	High lat. type
1158	36	2	P	Hæmorrhage one hour before admission	—	+	Cæsarean section	A	SB	Marginal
1259	30	7	FT	Fairly good	—	+	Cæsarean section	A	A	
1811	26	3	P	Good	—	+	Podalic version	A	A	
1866	26	8	FT	Fair, slight bleeding	+	—	Cæsarean section	A	A	Spinal anaesthesia. Little blood lost.
2692	21	3	FT	Fair	—	+	Cæsarean section	A	A	Marginal
2808	35	6	FT	Fairly good	—	+	Spontaneous del. pads and binder	A	SB	Marginal. Admitted advanced in labour.
2813	30	2	FT	Fair, vagina packed outside	—	+	Spontaneous delivery	A	A	Admitted advanced in labour
3820	36	8	P	Fair, moderate hæmorrhage outside	—	+	Breech, foot brought down	A	D	
3882	22	1	FT	Exsanguinated. Breech presenting	—	+	Memb. rupt. foot brought down	A	SB	Marginal. Blood transfusion
3885	34	4	FT	Severe hæmorrhage	—	+	Packed podalic version	A	SB	Marginal

Reg. No.	Age.	Parity.	Maturity.	Condition on Admission.	Type.		Treatment	Result.		Remarks.
					Cnt.	Lat.		M.	Ch.	
3916	25	2	P	Poor	—	+	Memb. rupt., podalic version	A	SB	Marginal
3933	36	8	FT	Marked pallor	—	+	Pads and binder, pituitrin	A	A	
2165	22	2	FT	Fair	—	+	Pads and binder, spontaneous delivery	A	A	High lat. placenta prævia
3772	22	3	P	Fairly good	—	+	Spontaneous delivery	A	A	Admitted advanced in labour

B. Cases sent in as Emergencies.

48	27	2	P	Fair	+	—	Cæsarean section	A	A	
82	33	3	P	Fair	+	—	Packing and version	A	SB	Blood transfusion
122	23	1	P	Poor, collapsed	+	—	Packed, spontaneous delivery	A	SB	
218	35	7	FT	Good	—	+	Podalic version	A	SB	Marginal
302	37	6	P	Poor, severe bleeding. Packed outside	+	—	Packing and podalic version	D	SB	Puer. septicæmia (Plac. manually removed)
361	39	8	FT	Fair, moderate bleeding	+	—	Podalic version	A	SB	
578	27	3	P	Fairly good	—	+	Packing	A	A & A	Large placenta
584	18	1	FT	Good	—	+	Spontaneous delivery	A	A	G.C. and W.R.+
608	32	2	P	Fairly good	—	+	Memb. rupt. Pituit. Manual delivery	A	SB	High lat. type
628	27	3	P	Good	—	+	Packing and version	A	SB	Marginal
667	41	8	P	Collapsed. Exsanguinated	+	—	Packing and intravenous gum acacia, etc.	D	—	Died 2½ hours after admission
675	31	5	FT	Fairly good	+	—	Podalic version	A	SB	
706	39	10	P	Fairly good	—	+	Podalic version	A	SB	
752	33	2	P	Good	+	—	Cæsarean section	A	A	Puer. pyrexia. Good recovery
936	29	5	FT	Good	—	+	Spontaneous delivery	A	SB	High lat. type with prol. cord

Reg. No.	Age.	Parity.	Maturity.	Condition on Admission.	Type.		Treatment.	Result.		Remarks.
					Cnt.	Lat.		M.	Ch.	
962	29	3	FT	Poor	—	+	Podalic version	A	A	Marginal
1017	29	3	P	Fairly good	+	—	Cæsarean section	A	A	
1022	44	14	P	Fairly good	+	—	Cæsarean section	A	SB	
1285	42	11	FT	Good	—	+	Spontaneous delivery	A	SB	High lat. type, cord prolapsed Hæmolytic strep- tococcal septicæmia
1351	26	4	P	Collapsed	+	—	Podalic version	D	A	
1463	29	3	P	Poor	+	—	Podalic version	A	SB	
1477	37	4	FT	Fair	—	+	Spontaneous delivery	A	A	High lat. type
1508	31	5	FT	Fair	—	+	Spontaneous delivery	A	A	High lat. type
1541	40	11	P	Collapsed	—	+	Podalic version	A	SB	Marginal
1599	38	8	FT	Collapsed	+	—	Cæsarean section and sterilization	A	A	Spinal anæsthetic
1623	30	1	FT	Fairly good	+	—	Cæsarean section	A	A	
1624	—	3	FT	Fairly good	+	—	Podalic version	A	SB	
1640	24	2	P	Fair, slight hæmorrhage	—	+	Foot brought down	A	SB	
1808	34	2	FT	Fairly good	+	—	Packed. Cæsarean section	A	A	
1829	38	8	P	Poor, severe hæmorrhage	+	—	Podalic version	A	SB	
1848	32	2	P	Fair, moderate hæmorrhage	+	—	Cæsarean section	A	D	
1927	34	2	FT	Collapsed, severe hæmor- rhage	+	—	Packed. Later, podalic version	A	SB	
2048	42	10	P	Poor, slight bleeding	—	+	Podalic version	A	A	
2050	28	5	FT	Severe bleeding	+	—	Cæsarean section	A	A	Severe mitral stenosis
2113	34	6	FT	Good	—	+	Spontaneous delivery	A	A	
2188	40	12	P	Fair	—	+	Cæsarean section	A	SB	
2190	39	6	FT	Poor, severe bleeding	+	—	Cæsarean section	A	A	Hydramnios present. Patient sterilized

Reg. No.	Age.	Parity.	Maturity.	Condition on Admission.	Type.		Treatment.	Result.		Remarks.
					Cnt.	Lat.		M.	Ch.	
2380	34	5	FT	Poor, severe bleeding. Packed	+	—	Podalic version	A	SB	
2382	36	2	FT	Good	—	+	Spontaneous delivery	A	A	High lat. type
2600	29	3	FT	Fair	+	—	Packing and cæsarean section	A	A	
2602	28	2	P	Poor, collapsed	+	—	Blood transfusion, version	A	SB	
2612	36	9	FT	Poor, collapsed. Packed	+	—	Cæsarean section	A	A	Puer. sepsis. Good recovery
2700	34	5	FT	Good	—	+	Memb. rupt., Pituitrin, spontaneous delivery	A	A	Marginal
2747	32	3	P	Fair	—	+	Cæsarean section	A	A.D	Marginal
2794	33	1	P	Fair	—	+	Version	A	A.D	Marginal P.P.H.
2800	34	1	P	Fairly good	+	—	Cæsarean section	A	A	
2817	33	4	P	Fair. Packed	+	—	Cæsarean section	A	A.D	
2868	30	6	P	Exsanguinated, moribund	+	—	Packing and intravenous gum acacia	D	—	Died 2 hours after admission undelivered.
2971	24	2	FT	Poor	—	+	Podalic version	A	SB	Marginal. Developed pelvic cellulitis. Good recovery
3014	31	6	FT	Good	—	+	Spontaneous delivery	A	A	High lat. type
3085	21	1	P	Fairly good	—	+	Podalic version	A	SB	Marginal
3137	43	10	P	Good	+	—	Cæsarean section	A	A.D	Feeble child
3191	38	12	P	Severe bleeding	—	+	Podalic version	A	SB	Marginal
3239	36	2	P	Good	—	+	Cæsarean section	A	A.D	Marginal
3241	40	5	FT	Fairly good	—	+	Cæsarean section	D	A	Collapsed and died day after operation No P.M.
3449	32	3	FT	Severe bleeding	—	+	Packing	A	A	Marginal
3716	22	4	P	Fair	—	+	Podalic version	A	SB	
3743	25	3	P	Fairly good	—	+	Memb. rupt., spontaneous delivery	A	A	Marginal. Puer. sepsis. Good recovery
3793	38	1	FT	Slight bleeding	—	+	Packing and forceps	A	SB	

Reg. No.	Age.	Parity.	Maturity.	Condition on Admission.	Type.		Treatment.	Result.		Remarks.
					Cnt.	Lat.		M.	Ch.	
3823	28	2	FT	Poor	—	+	Packing	A	SB	Puer. sepsis Blood transfusion in puerperium
3941	32	3	P	Collapsed, severe hæmorrhage	+	—	Blood transfusion, version	A	SB	
4100	26	6	P	Poor, severe bleeding	—	+	Memb. rupt., packing	A	SB	Marginal
4199	29	1	FT	Good, packed	+	—	Cæsarean section	A	A	
4423	33	7	P	Poor, severe bleeding	+	—	Podalic version, blood transfusion	A	SB	

C.—Not under Ante-natal Supervision.

857	36	1	FT	Fair	+	—	Cæsarean section (spinal anæsthesia)	A	A	Double phlegmasia alba dolens
239	29	3	P	Good	—	+	Packing	A	SB	Marginal
1036	40	8	P	Fairly good	+	—	Cæsarean section (Spinal Anæsthetic)	A	A.D	
1128	41	4	P	Fair	+	—	Cæsarean (Spinal Anæsthetic)	A	A.D A.D	
1509	26	5	FT	Poor, severe bleeding	—	+	Spontaneous delivery on adm.	A	SB	
1938	27	3	P	Fair	—	+	Cæsarean section	D	AD	Puer. sepsis. Broncho- pneumonia
1930	30	5	P	Fairly good	+	—	Cæsarean section	A	D	
2431	34	4	FT	Good, slight bleeding	—	+	Spontaneous	A	A	High lat. type
2672	31	6	FT	Severe bleeding, collapsed	—	+	Version	D	SB	
2950	29	7	P	Fairly good	—	+	Packing, spontaneous delivery	A	SB	Marginal
3495	34	6	P	Fairly good	—	+	Memb. rupt., spontaneous delivery	A	A	Marginal puer. pyrexia. Good recovery
3911	24	4	FT	Good	—	+	Spontaneous delivery	A	A	High lat. type

ACCIDENTAL HÆMORRHAGE.

A.—Cases under Ante-natal Supervision.

Reg. No.	Age.	Parity.	Maturity.	Condition on Admission.	Type.		Treatment.	Result.		Remarks.
					Apparent.	Concealed.		M.	Ch.	
17	27	4	P	Moderate bleeding	+	+	Memb. rupt., spontaneous delivery	A	SB	
286	24	2	FT	Good	+	—	Med. ind., spontaneous delivery	A	A	Mild case
406	39	4	P	Severe shock	—	+	Intravenous gum acacia sedatives, packing	D	—	
510	34	6	FT	Slight bleeding	+	—	Spontaneous delivery	A	A	? Excessive shows
711	27	2	FT	Slight bleeding	+	—	Spontaneous delivery	A	A	
731	37	1	FT	Good, slight bleeding	+	—	Forceps delivery	A	SB	
906	35	8	FT	Fairly good, moderate bleeding	+	—	Memb. rupt. pituitrin spontaneous delivery	A	A	Mild case
931	34	1	FT	Good, slight bleeding	+	—	Forceps	A	A.D	Feeble child
1227	23	2	P	Profuse bleeding	+	—	Spontaneous breech delivery	A	SB	
1684	35	5	P	Slight bleeding	+	—	Spontaneous breech delivery	A	SB	
1904	34	6	P	Slight bleeding	+	—	Spontaneous delivery	A	SB	Previous operation for left-sided pyosalpiux
2325	32	9	P	Good	+	—	Rest	A	—	Irreg. dismissal undelivered. W.R. + +
2688	40	5	FT	Very slight bleeding	+	—	Spontaneous delivery	A	A	
3197	22	2	FT	Slight pain and bleeding	+	—	Spontaneous delivery	A	A	? Excessive "shows"
3313	32	6	FT	Slight pain and bleeding	+	—	Spontaneous delivery	A	A	? Excessive "shows"

Reg. No.	Age.	Parity.	Maturity	Condition on Admission.	Type.		Treatment.	Result.		Remarks
					Apparent.	Concealed.		M.	Ch.	
3701	38	7	P	Pain, bleeding and shock	+	+	Rupture of membranes, spontaneous delivery	A	SB	Large retroplacental hæmatoma (1½ lbs.)

B.—Cases sent in as Emergencies.

41	39	10	P	Fair, moderate bleeding	+	—	Memb. rupt., spontaneous delivery	A	SB	Mild case
76	19	2	FT	Good, slight bleeding	+	—	Spontaneous delivery	A	A	
99	24	2	FT	Collapsed, severe bleeding	+	—	Pituitrin and packing	A	SB	
164	26	5	FT	Poor, severe bleeding	—	+	Morphia and pituitrin	A	SB	Large retroplacental hæmatoma (1¼ lbs.)
217	33	5	P	Fair, moderate bleeding	+	+	Morphia and pituitrin	A	A	
310	40	13	P	Fair	+	+	Morphia and pituitrin	D	—	Severe shock manifested shortly before death
355	29	3	FT	Fairly good, slight bleeding	+	—	Spontaneous delivery, plac. manually removed	A	SB	Puer. pyrexia. Good recovery
435	36	5	P	Fairly good, slight bleeding	+	—	Spontaneous delivery	A	A	Fecble child
437	39	9	FT	Poor	+	+	Intravenous gum acacia morphia, pituitrin	A	SB	Large retroplacental hæmatoma
445	22	1	FT	Good, slight bleeding	+	—	Spontaneous delivery	A	A	Mild case
535	37	10	FT	Moribund	—	+	Shock treated	D	—	Died 1 hour after admission
538	32	10	?	Collapsed, severe shock	—	+	Intravenous saline	D	—	Died 30 minutes after admission
782	38	3	FT	Poor, severe bleeding	+	+	Memb. rupt., spontaneous delivery	A	A	Small retroplacental clot

Reg. No.	Age.	Parity.	Maturity.	Condition on Admission.	Type.		Treatment.	Result.		Remarks.
					Apparent.	Concealed.		M.	Ch.	
789	24	2	FT	Slight bleeding	+	—	Packing and pituitrin	A	A	Feeble child
819	24	3	FT	Good	+	—	Spontaneous delivery	A	A	Mild case
856	37	10	FT	Fairly good, slight bleeding	+	—	Memb. rupt., morphia	A	A	.
970	34	5	P	Severe shock	—	+	Sedatives, spontaneous delivery	A	SB	
1025	43	12	FT	Good, slight bleeding	+	+	Cæsarean section	A	SB	Intra-mural uterine hæmorrhages
1027	22	1	FT	Fairly good	+	—	Pituitrin spontaneous delivery	A	A	
1041	28	2	FT	Slight bleeding	+	—	Sedatives and rest	A	—	Irregular dismissal
1162	33	5	FT	Slight bleeding	+	—	Spontaneous delivery	A	A	Mild case
1228	34	1	P	Slight bleeding	+	—	Spontaneous delivery	A	A	.
1350	38	10	P	Severe shock, slight bleeding	+	+	Sedatives, spontaneous delivery	A	SB	
1430	26	4	FT	Slight bleeding	+	—	Spontaneous delivery	A	A	Slight apparent acc. hæm.
1436	44	9	P	Severe shock	—	+	Memb. ruptured, spontaneous delivery	A	SB	Large retroplacental hæmatoma (1 $\frac{3}{4}$ lbs.)
1560	34	5	P	Pain and bleeding	+	+	Spontaneous delivery	A	A	
1584	18	1	P	Collapsed, severe shock	—	+	Memb. ruptured, spontaneous delivery	A	SB	Large retroplacental hæmatoma
1602	23	2	P	Brisk hæmorrhage	+	—	Sedatives, spontaneous delivery	A	AD	
1626	19	3	P	Collapsed, marked tender- ness and rigidity of abdomen	+	—	Memb. rupt., spontaneous delivery	A	SB	Retroplacental hæmatoma
1649	—	—	P	Severe pain, shock	+	—	Sedatives, spontaneous delivery	A	SB	

Reg. No.	Age.	Purity.	Maturity.	Condition on Admission.	Type.		Treatment.	Result.		Remarks.
					Apparent.	Concealed.		M.	Ch.	
1716	36	8	FT	Profuse bleeding	+	—	Spontaneous delivery	A	A	
1725	22	3	P	Slight bleeding	+	+	Sedatives, spontaneous delivery	A	SB	
1839	35	3	P	No bleeding	—	+	Sedatives, spontaneous delivery	A	SB	Chronic nephritis
2003	—	9	FT	Slight bleeding	+	—	Spontaneous delivery	A	A	? Excessive shows
2157	44	13	P	Pain, shock and profuse bleeding	+	+	Memb. rupt., packed, pituitrin, spontaneous delivery	A	SB	Large retroplacental hæmorrhage
2182	33	2	—	Pain, no bleeding	—	+	Complete hysterectomy	A	SB	Anuria on admission. A typical case
2215	41	4	FT	Slight bleeding	+	—	Spontaneous delivery	A	SB	
2243	36	1	FT	Slight bleeding, packed	+	+	Version to footling	A	SB	Retroplacental hæmatoma
2346	40	12	FT	Profuse bleeding, collapsed	+	+	Spontaneous delivery	A	SB	
2353	31	3	P	Collapsed	+	—	Spontaneous delivery	A	SB	
2451	35	4	P	Profuse bleeding	+	—	Memb. rupt., spontaneous delivery	A	SB	
2456	35	9	P	Severe shock	—	+	Shock treated, sedatives	D	—	Died 40 minutes after admission to Hospital
2488	37	3	P	Poor, toxic	+	+	Sedatives, memb. rupt., blood transfusion	D	—	Pre-eclamptic toxæmia and acc. hæm.
2501	39	11	P	Collapsed. Profuse hæm.	+	+	Memb. rupt., pituitrin and morphia, spontaneous delivery	A	SB	
2617	33	7	P	Shocked and very toxic	+	+	Pituitrin and morphia, spontaneous delivery	D	SB	Puerperal insanity. Suicide on 4th day
2646	42	10	FT	Shocked and exsanguinated	+	+	Packed, pituitrin and morphia, spontaneous delivery	D	SB	

Reg. No.	Age.	Parity.	Maturity.	Condition on Admission.	Type.		Treatment.	Results.		Remarks.
					Apparent.	Concealed.		M.	Ch.	
2797	32	5	FT	Marked shock, profuse bleeding. vagina packed	+	+	Memb. rupt., forceps delivery	D	SB	Died 3 hours after admission
2815	39	7	P	Shocked and very toxic	+	+	Memb. rupt. packed, Pituitrin, morphia	D	—	P.M. Large retro- placental clot. Acute endocarditis (Result of toxæmia)
2891	30	6	P	Slight bleeding	+	+	Memb. rupt., binder and pads	A	A	Small retroplacental hæmatoma
2904	25	5	P	Slight bleeding	+	—	Spontaneous delivery	A	A.D	Mild case
3017	35	8	FT	Good. Vagina, packed outside	+	—	Spontaneous delivery	A	A	Mild case
3038	34	9	P	Fair, marked pallor	+	+	Spontaneous delivery	A	A.D	Large retroplacental clot. Feeble infant
3096	30	11	P	Collapsed, slight bleeding	+	—	Vagina packed, spontaneous delivery	A	A	Small retroplacental clot
3107	35	7	P	Profuse bleeding	+	—	Memb. rupt.	A	A.D	Very feeble child
3117	34	2	P	Poor, marked pallor	+	+	Spontaneous delivery	A	SB	Small retroplacental clot
3119	30	4	P	Marked pallor and shock	—	+	Sedatives	A	SB	Large retroplacental clot (1½ lbs.)
3132	26	4	FT	Good	+	—	Memb. rupt.	A	A & A	Mild case
3200	42	14	FT	Severe shock	—	+	Memb. rupt. forceps delivery	A	SB	Large retroplacental clot
3228	23	3	FT	Severe shock, Collapsed	—	+	Pituitrin. and sedatives, spontaneous delivery	A	SB	Large retroplacental clot
3262	19	2	P	Fairly good	—	+	Sedatives	A	SB	Large retroplacental clot (1 lb.)
3327	41	3	FT	Slight bleeding	+	—	Memb. rupt.	A	A	Mild case
3374	26	7	FT	Fairly good	—	+	Memb. rupt. spontaneous delivery	A	SB	Large retroplacental clot

Reg. No.	Age.	Purity.	Maturity.	Condition on Admission.	Type.		Treatment.	Result.		Remarks.
					Apparent.	Concealed.		M.	Ch.	
3399	28	5	FT	Albuminuria and slight vaginal bleeding	+	+	Sedatives and pituitrin	A	SB	Large retroplacental clot (1 $\frac{3}{4}$ lbs.)
3428	22	2	P	Collapsed	+	+	Sedatives, memb. rupt., pituitrin	A	SB	Delivered after version to breech
3567	37	7	FT	Severe bleeding	+	+	Memb. rupt., sedatives	A	SB	Large retroplacental clot (2 $\frac{1}{2}$ lbs.)
3610	26	3	P	Marked pallor and bleeding	+	—	Memb. rupt., spontaneous delivery	A	A.D	Small retroplacental clot and lateral plac. prævia
3638	37	10	P	Slight bleeding	+	—	Sedatives, spontaneous delivery	A	A	? Excessive shows
3781	26	4	P	Marked pallor, slight bleeding	+	—	Sedatives, spontaneous delivery	A	SB	Apparent acc. hæm. and lat. plac. prævia
3785	25	3	FT	Slight bleeding	+	+	Spontaneous delivery	A	A	Retroplacental clot ($\frac{3}{4}$ lb.)
3812	26	5	FT	Slight bleeding	+	—	Spontaneous delivery	A	SB	Prol. cord
3855	39	4	P	Severe album.	+	+	Spontaneous delivery	A	SB	3 weeks in A.N. wards. Albuminuria
3910	32	9	P	Severe pain and shock	—	+	Sedatives and pituitrin	A	SB	
4029	37	12	FT	Slight shock, abdom. tenderness	—	+	Cæsarean section and sterilization	A	SE	
4047	37	5	P	Severe shock	—	+	Sedatives	A	SB & SB	Retroplacental clot (2 lbs)
4071	26	3	P	Shock and bleeding	+	+	Sedatives and pituitrin memb. rupt., breech delivery	A	SB	
4079	34	8	P	Abdom. tenderness, shock	—	+	Sedatives, pituitrin, forceps delivery	A	SB	
4103	34	8	P	Profuse bleeding	+	—	Memb. rupt., vagina packed	A	A	

Reg. No.	Age.	Parity.	Maturity.	Condition on Admission.	Type.		Treatment.	Result.		Remarks.
					Apparent.	Concealed.		M.	Ch.	
4118	33	10	P	Good	—	+	Manual extraction	A	SB	Anencephalic monster
4196	28	4	P	Severe bleeding	+	+	Sedatives	A	SB	Retroplacental clot ($\frac{1}{2}$ lb.)
4388	36	9	P	Marked pallor, collapsed	+	+	Sedatives	A	SB	Retroplacental clot ($1\frac{1}{4}$ lbs.). Marked albuminuria
4468	32	7	P	Shocked	—	+	Sedatives	A	SB	Retroplacental clot ($1\frac{3}{4}$ lbs.)
<i>C.—Cases not under Ante-natal Supervision.</i>										
197	25	1	P	Good, slight bleeding	+	—	Rest and chlor. exam.	A	—	No further bleeding for 3 weeks. No placenta felt
911	31	6	P	Fairly good, slight bleeding	+	—	Morphia, spontaneous delivery	A	A.D	
925	34	10	P	Collapsed	+	+	Intravenous gum acacia, memb. rupt., spontaneous delivery	A	SB	Large retroplacental clot
968	24	3	FT	Good	+	—	Sedatives, spontaneous delivery	A	A	Mild case
1940	23	2	P	Fairly good	+	—	Spontaneous delivery	A	A.D	Feeble infant. Mild case
2267	30	2	P	Collapsed, severe shock	+	+	Morphia and pituitrin, spontaneous delivery	A	SB	Large retroplacental clot
2586	29	6	P	Good	+	—	Spontaneous delivery	A	A	Feeble child. Mild case
2756	19	1	P	Good	+	—	Spontaneous delivery	A	A	Feeble child. Mild case
2968	37	14	FT	Slight bleeding	+	—	Spontaneous delivery	A	A	
3158	37	10	P	Slight bleeding	+	—	Spontaneous delivery	A	A	
3160	21	3	FT	Slight bleeding	+	—	Spontaneous delivery	A	A	
3667	35	8	P	Marked shock, severe toxæmia	—	+	Memb. rupt., spontaneous delivery	D	SB	P.M. acid hæm. and toxic jaundice
3997	26	5	P	Severe shock, collapsed	—	+	Sedatives, spontaneous delivery	A	SB	Retroplacental blood clot ($1\frac{1}{2}$ lbs.)
4194	23	3	P	Good	+	—	Rest	A	—	1 week in A.N. Ward
4371	22	3	P	Profuse bleeding	+	+	Spontaneous delivery	A	A.D	

TOXÆMIA OF PREGNANCY.

During the year 1930 there were 448 cases of toxæmia of pregnancy of which records are available. Owing to lack of uniformity in the nomenclature employed by different members of the staff, it has been necessary to confine the subdivision of these cases to three classes, viz. :—(1) albuminuria, pre-eclamptic toxæmia, and nephritic toxæmia ; (2) eclampsia ; (3) hyperemesis (including late persistent vomiting). Cases of transient albuminuria during labour or immediately following it have been excluded. Dividing the cases in this way, there were 279 cases of the first class (albuminuria, pre-eclamptic toxæmia, and nephritic toxæmia), 59 of the second class (eclampsia), and 110 of the third class (hyperemesis and late vomiting).

Amongst the 1,790 cases which had been under ante-natal supervision there were 173 toxæmia cases (134 in class 1, 7 in class 2, and 32 in class 3). There were thus 7.4 per cent. of the cases in Category A which were toxæmic (class 1), and 3.9 per cent. which were eclamptic (class 2).

Of the 1,934 cases which were sent in as emergencies there were 250 which were toxæmic (126 in class 1, 49 in class 2, and 75 in class 3). There were thus 6.5 per cent. of the cases in Category B which were toxæmic (class 1) and 2.5 per cent. which were eclamptic (class 2).

Of the 745 cases which were admitted without previous supervision there were 25 which were toxæmic (19 in class 1, 3 in class 2, and 3 in class 3). There were thus 2.5 per cent. of the cases in Category C which were toxæmic (class 1) and 40 per cent. which were eclamptic (class 2).

There were 22 maternal deaths out of the total 448 toxæmic cases. The incidence in the various groups is shown in the following table :—

Table showing the Maternal Deaths from the various forms of Toxæmia in each of the three Categories A, B, and C.

Type of Toxæmia.	Categ. A.	Categ. B.	Categ. C.	Total Deaths.
1. Albuminuria, pre-eclamptic toxæmia and nephritic toxæmia - -	4	5	—	9
2. Eclampsia - - - -	1	7	—	8
3. Hyperemesis gravidarum (including late persistent vomiting) - -	—	4	1	5
Total deaths - -	5	16	1	22

ALBUMINURIA, PRE-ECLAMPTIC TOXÆMIA, AND NEPHRITIC TOXÆMIA.

A.—Cases under Ante-natal Supervision.

Reg. No.	Age.	Parity.	Stage of Pregnancy.	Mode of Delivery.	Result.					Remarks.	
					Mother.		Child.				
					L.	D.	L.	D.	S.B.		
18	31	2	P	Spontaneous	-	+	-	+	-	-	Mild albuminuria.
107	28	1	FT	Undelivered	-	+	-	-	-	-	Chronic nephritis. Esbach, 2 parts. B.P. 130/100.
177	26	1	P	Induction (Bougies)	-	-	+	-	-	+	Severe alb. Esb. 2-6 parts.
299	20	1	FT	Spontaneous	-	+	-	+	-	-	Esb. 1-2 parts. B.P. 130/100.
360	24	3	P	Undelivered	-	+	-	-	-	-	Marked œdema. Sl. albuminuria.
378	19	1	FT	Spontaneous	-	+	-	+	-	-	Esb., 1 part. Pre- eclamptic symptoms.
386	29	2	6	Undelivered	-	+	-	-	-	-	Mild case.
632	35	3	P	Spontaneous	-	+	-	+	-	-	Mild albuminuria. œdema.
639	19	2	FT	Spontaneous	-	+	-	+	-	-	Mild albuminuria. œdema.
672	21	1	FT	Spontaneous	-	+	-	+	-	-	Esbach, ½ part. Mild case.
709	20	1	38 wks.	Surgical induction, spontaneous.	-	+	-	+	-	-	Esbach, 1-4 parts. B.P. 140/80.
717	24	1	FT	Spontaneous	-	+	-	+	-	-	Esbach, ½-1 part.
741	27	2	8	Undelivered	-	+	-	-	-	-	Alb. trace. Mild case.
745	38	3	32 wks.	Surgical induction (bougies)	-	-	+	-	-	+	Esbach, 1-2 parts. B.P., 190/140. Chronic nephritis. Previous stillbirths
755	21	2	8	Undelivered	-	+	-	-	-	-	Esbach, ½ part. Mild case
777	20	1	FT	Spontaneous medical induction	-	+	-	+	-	-	Esbach, 3-6 parts. B.P. 180/130. Pre-eclamp- tic symptoms
794	35	12	FT	Spontaneous	-	+	-	++	-	-	Esbach, ½ part. Pre- eclamptic symptoms
884	20	1	FT	Spontaneous	-	+	-	+	-	-	Esbach, 1 part. Pyelitis
887	30	3	P	Spontaneous	-	+	-	-	-	+	Esbach, 2-3 parts
890	31	5	FT	Spontaneous	-	+	-	+	-	-	Alb. trace. Chronic nephritis
896	22	1	8	Undelivered	-	+	-	-	-	-	Esbach, ½ part. Mild case
899	22	1	8	Undelivered	-	+	-	-	-	-	Esbach, 1-3 parts. B.P., 130/100. Irregular dism.
922	23	1	FT	Spontaneous	-	+	-	+	-	-	œdema. B.P. 70/108
940	34	9	26 wks.	Surgical induction (bougies)	-	+	-	+	-	-	Esbach, 6-14 parts. B.P., 160/90. Chronic nephritis. 5 previous stillbirths

Reg. No.	Age.	Parity.	Stage of Pregnancy.	Mode of Delivery.	Result.					Remarks.
					Mother.		Child.			
					L.	D.	L.	D.	S.B.	
1010	20	1	7	Undelivered -	+	-	-	-	-	Alb. trace. Mild case
1081	23	1	FT	Medical induction, spontaneous	+	-	+	-	-	Pre-eclamptic symp- toms. B.P., 145/80
1104	34	1	F	Undelivered	+	-	-	-	-	B.P. 120/80. Esb. 1 part
1116	20	1	FT	Spontaneous -	+	-	+	-	-	Esb. 1 part. B.P. 120/70
1131	19	1	FT	Medical induction, spontaneous	+	-	+	-	-	Esbach, 3 parts. B.P. 190/118
1161	40	10	8	Undelivered -	+	-	-	-	-	Trace alb. Mild case
1189	28	2	FT	Forceps and manual	+	-	++	-	-	Esb. 1 part. B.P. 140/100
1190	31	1	FT	Spontaneous -	+	-	+	-	-	Trace alb. B.P., 120/70. Mild case
1192	24	2	FT	Spontaneous -	-	+	+	-	-	Esb. 1 part. Mild case
1250	41	10	FT	Spontaneous -	+	-	+	-	-	Esb. 1-6 parts. B.P. 170/100. Pre-eclamp- tic tox. 1 day in hospital
1253	26	4	FT	Undelivered -	+	-	-	-	-	Irregular dism.
1275	22	1	FT	Spontaneous -	+	-	+	-	-	Esb. 2-14 parts. B.P. 150/100. Pre-eclam. tox
1288	26	4	8	Undelivered -	+	-	-	-	-	Esb. 1-5 parts. B.P. 125/80
1289	31	5	FT	Medical induction, spontaneous	+	-	+	-	-	Esb. ½-4 parts. B.P. 120/70
1290	27	1	8	Spontaneous -	+	-	+	-	-	Esb. 2-8 parts. B.P. 180/110. Pre-eclam. tox
1328	22	1	8	Undelivered -	+	-	-	-	-	Trace alb. Mild case. B.P. 140/100
1330	19	1	FT	Spontaneous -	+	-	+	-	-	Esb. ½-4 parts
1364	25	1	FT	Spontaneous -	+	-	+	-	-	Esb. ½-4 parts
1373	21	3	7	Undelivered -	+	-	-	-	-	Esb. 2 parts. B.P. 120/80. Irregular dism.
1481	23	1	7	Undelivered -	+	-	-	-	-	Esb. 1-7 parts. B.P. 160/100
1482	21	1	FT	Spontaneous -	+	-	-	-	+	Esb. 1-4 parts. B.P. 160/80
1543	38	8	7	Spontaneous -	+	-	+	-	-	Esb. 1-6 parts. B.P. 130/90
1546	21	2	FT	Spontaneous -	+	-	+	-	-	Esb. ½ part. B.P. 130/80
1727	23	2	8	Manual -	+	-	+	-	-	Alb., 1-5 parts. B.P. 180/120
1728	17	1	FT	Spontaneous -	+	-	+	-	-	Esb. 1-5 parts. B.P. 140/90
1763	25	6	5	Undelivered -	+	-	-	-	-	Alb. trace. B.P. 150/80. Mild case

Reg. No.	Age.	Parity.	Stage of Pregnancy.	Mode of Delivery.	Result.					Remarks.	
					Mother.		Child.				
					L.	D.	L.	D.	S.B.		
1801	29	7	8	Spontaneous	-	+	-	+	-	-	Esb. 1-3 parts. B.P. 140/100
1834	23	1	FT	Spontaneous	-	+	-	+	-	-	Alb. trace. B.P. 90/60. Mild case
1873	23	1	8	Spontaneous	-	+	-	+	-	-	Alb. trace. Mild case
1880	41	3	FT	Spontaneous	-	+	-	+	-	-	Alb. trace. Mild case. B.P., 120/80
1993	26	1	FT	Forceps	-	+	-	+	-	-	Esb. $\frac{1}{2}$ part. B.P. 120/80. Pre-eclam. tox.
1996	23	1	FT	Spontaneous	-	+	-	+	-	-	Alb. trace. B.P. 118/60. Mild case
2140	26	4	FT	Spontaneous	-	+	-	++	-	-	Alb. trace. B.P. 120/90. Mild case
2143	33	1	7	Undelivered	-	+	-	-	-	-	Trace alb. 3 days in hospital. Irregular dism.
2159	41	6	FT	Spontaneous	-	+	-	+	-	-	Esb. $\frac{1}{2}$ -2 parts. B.P. 160/90. Chronic nephritis
2160	26	1	FT	Surgical induction (bougies)		+	-	+	-	-	Esb. 1-3 parts. B.P. 150/110. Pre-eclam. tox.
2181	21	1	FT	Spontaneous	-	+	-	+	-	-	Alb. trace. Mild case
2183	33	10	5	Induction of abortion (tents)		+	-	-	-	-	Esb. 2-4 parts. B.P. 190/150
2211	25	1	FT	Manual	-	+	-	+	-	-	Esb. 2 parts. B.P. 160/100
2324	23	4	FT	Spontaneous	-	+	-	+	-	-	Esb. 1 part. B.P. 170/120. Pre-eclam. tox.
2541	30	1	FT	Forceps	-	+	-	+	-	-	Alb. trace. B.P. 140/100. Mild case
2579	28	4	8	Caes. section	-	+	-	+	-	-	Esb. $\frac{1}{2}$ -3 parts. B.P. 160/90. Pre-eclam. tox.
2643	26	3	FT	Spontaneous	-	+	-	+	-	-	Esb. $\frac{1}{2}$ -2 parts. B.P. 120/70
2677	21	2	FT	Spontaneous	-	+	-	+	-	-	Alb. trace. B.P. 130/100. Mild case
2689	25	1	6	Undelivered	-	+	-	-	-	-	Alb. trace. B.P. 130/80. Mild case
2696	27	3	8	Undelivered	-	+	-	-	-	-	Esb. $\frac{1}{2}$ part. B.P. 110/90
2704	28	5	FT	Spontaneous	-	+	-	+	-	-	Esb. $\frac{1}{2}$ -1 part. B.P. 130/90
2734	32	2	8 $\frac{1}{2}$	Undelivered	-	+	-	-	-	-	Alb. trace. B.P. 130/70. Mild case
2776	26	1	FT	Forceps	-	+	-	+	-	-	Esb. $\frac{1}{2}$ part. Oedema
2811	29	1	FT	Manual extractions (2) and spontaneous (1)		+	-	+	+	-	Alb. trace. B.P. 130/80. Mild case

Reg. No.	Age.	Parity.	Stage of Pregnancy.	Mode of Delivery.	Result.					Remarks.
					Mother.		Child.			
					L.	D.	L.	D.	S.B.	
2849	28	1	FT	Forceps - -	+	-	+	-	-	Esb. $\frac{1}{2}$ part
2883	22	1	FT	Spontaneous -	+	-	+	-	-	Esb. $\frac{1}{2}$ part. B.P. 140/90
2885	39	8	8 $\frac{1}{2}$	Undelivered -	+	-	-	-	-	Alb. trace. B.P. 120/80. Mild case
2911	27	4	FT	Medical induction, spontaneous	+	-	+	-	-	Esb. 1-7 parts. B.P. 150/90. Pre-eclam. tox.
2937	26	3	FT	Spontaneous -	+	-	+	-	-	Alb. trace. Mild case
2984	29	1	FT	Spontaneous -	+	-	+	-	-	Esb. $\frac{1}{2}$ -3 parts. B.P. 160/110. Pre-eclam, tox.
3000	30	4	FT	Spontaneous -	+	-	+	-	-	Esb. $\frac{1}{2}$ part. B.P. 160/90. M a r k e d œdema
3005	31	5	FT	Spontaneous (breech) -	+	-	+	-	-	Esb. 1-3 parts. B.P. 110/70
3012	23	1	FT	Spontaneous -	+	-	+	-	-	Alb. trace., Mild case. B.P. 130/90
3025	23	1	P	Manual extraction	+	-	+	-	-	Alb. trace. Pre-eclam. tox.
3026	29	1	FT	Forceps - -	+	-	+	-	-	Pre-eclam. tox. Esb. $\frac{1}{2}$ part. B.P., 130/90
3072	20	1	P	Surgical induction, spontaneous	+	-	+	-	-	Esb. 1-4 parts. B.P. 150/100. Severe pre-eclam. tox.
3073	37	5	S	Spontaneous -	+	-	+	-	-	Pre-eclam. tox. Alb. trace. B.P., 120/70
3074	33	3	P	Surgical induction, spontaneous (bougies)	+	-	+	-	-	Esb. 1-12 parts. B.P. 190/40
3079	33	6	8 $\frac{1}{2}$	Undelivered -	+	-	-	-	-	Alb. trace. B.P. 145/95. Mild case
3081	26	4	7	Undelivered -	+	-	-	-	-	Alb. trace œdema. Mild case
3091	39	16	FT	Spontaneous -	+	-	+	-	-	Alb. trace. B.P. 180/110. Chronic nephritis
3092	35	1	P	Manual extraction of breech	+	-	-	-	+	Esb. $\frac{1}{2}$ part. B.P. 150/100
3148	22	4	FT	Spontaneous -	+	-	+	-	-	Alb. trace. B.P. 140/100. Mild case
3284	23	1	FT	Spontaneous -	+	-	+	-	-	Esb. 1-3 parts. B.P. 170/100. Pre-cclam. tox.
3392	40	8	FT	Undelivered -	-	+	-	-	-	Esb. 1-3 parts. B.P. 160/100. Chronic nephritic. Anæsthetic death.
3393	27	1	FT	Spontaneous -	+	-	+	-	-	Esb. 1-3 parts. B.P. 160/120. Pre-eclam. tox.

Reg. No.	Age.	Parity.	Stage of Pregnancy.	Mode of Delivery.	Result.					Remarks.
					Mother.		Child.			
					L.	D.	L.	D.	S.B.	
3465	26	4	7	Undelivered -	+	-	-	-	-	Alb. trace. B.P. 100/60. Chronic nephritis
3473	32	6	8½	Undelivered -	+	-	-	-	-	B.P. 140/80. Œdema
3540	23	3	8	Undelivered -	-	-	-	-	-	Alb. trace. B.P. 120/100
3544	35	4	FT	Forceps -	+	-	-	-	+	Esb. ½-1 part. B.P. 130/80
3546	27	4	FT	Spontaneous -	+	-	+	-	-	Esb. ½ part. B.P. 120/80. Chronic nephritis
3618	40	10	FT	Spontaneous -	+	-	+	-	-	Marked œdema
3652	38	10	P	Spontaneous -	+	-	+	-	-	Alb. trace. B.P. 120/60
3798	22	1	FT	Forceps -	+	-	+	-	-	Esb. 2-3 parts. B.P. 170/120. Pre-eclam. tox.
3800	25	1	FT	Spontaneous -	+	-	+	-	-	Alb. trace. Mild case
3818	23	3	FT	Spontaneous -	+	-	+	-	-	Alb. trace. Pre-eclam. tox. B.P. 140/90
3819	25	2	8	Undelivered -	+	-	-	-	-	Esb. 1-5 parts. B.P. 140/100
3831	19	1	8	Surgical induction, spontaneous (bougies)	+	-	+	-	-	Esb. 7 parts. B.P. 200/110. Pre-eclam. tox.
3843	25	3	FT	Spontaneous -	+	-	+	-	-	Alb. trace
3855	39	4	P	Spontaneous -	-	-	-	-	+	Marked œdema. Pre-eclam. tox.
3857	31	5	FT	Spontaneous -	+	-	+	-	-	Alb. trace. Mild case
3942	29	2	2½	Undelivered -	+	-	-	-	-	Persistent vomiting. Acetonuria. Toxic. Good recovery after 9 weeks
3951	40	4	6	Surgical induction, complete miscarriage	+	-	-	-	-	Esb. 2-6 parts. B.P. 160/95. Chronic nephritis
3990	31	1	7	Miscarriage -	+	-	-	-	-	Esb. 1-2 parts. B.P. 180/30. Pre-eclam. tox
3991	—	1	FT	Forceps -	+	-	+	-	-	B.P. 170/110. Alb. trace. Pre-eclam. tox.
4006	26	4	FT	Spontaneous -	+	-	+	-	-	Esb. 1-3 parts. B.P. 130/70
4016	29	3	—	Undelivered -	+	-	-	-	-	Alb. trace. B.P. 100/70. Mild case
4057	39	4	FT	Medieal induction, spontaneous	+	-	+	-	-	Alb. trace. Mild case. B.P. 130/90
4059	—	5	FT	Spontaneous -	+	-	+	-	-	—
4076	38	1	FT	Forceps -	+	-	+	-	-	Alb. trace. B.P. 120/80. Pre-eclam. tox.

Reg. No.	Age.	Parity.	Stage of Pregnancy.	Mode of Delivery.	Result.					Remarks.
					Mother.		Child.			
					L.	D.	L.	D.	S.B.	
4083	25	3	FT	Spontaneous -	+	-	+	-	-	Esb. 1-3 parts. B.P. 180/90
4105	24	1	8	Spontaneous -	+	-	+	-	-	Esb. 1-2 parts. B.P. 160/90
4153	32	3	FT	Spontaneous -	+	-	+	-	-	B.P. 115/80. Alb. trace. Mild case
4188	27	1	P	Manual -	+	-	-	-	-	—
4219	21	2	FT	Spontaneous -	+	-	+	-	-	Esb. 1-7 parts. B.P. 140/100
4231	22	1	FT	Forceps -	+	-	+	-	-	Alb. trace. Mild case
4284	26	1	FT	Spontaneous -	+	-	+	-	-	B.P. 140/90. Alb. trace. Mild case.
4317	32	3	FT	Spontaneous -	+	-	+	-	-	B.P. 160/110. Chronic nephritis. Alb. trace
4326	25	6	8	Undelivered -	+	-	-	-	-	Esb. 2-4 parts. B.P. 130/100. Chronic nephritis
4328	21	1	8	Surgical induction, spontaneous (bougies)	+	-	-	-	+	Esb. 2-9 parts. B.P. 210/140. Pre-eclam. tox.
4345	25	1	FT	Spontaneous -	+	-	+	-	-	Esb. 1 part. B.P. 150/70
4381	20	1	FT	Spontaneous -	+	-	+	-	-	Alb. trace. Mild case
4456	34	1	FT	Cæs. section -	+	-	+	-	-	Alb. trace. Mild case

B.—Cases sent in as Emergencies.

6	24	3	7	Spontaneous -	+	-	-	-	+	Esb. 1-6 parts. Œdema
23	24	1	FT	Spontaneous -	+	-	+	-	-	Esbach, 12 parts. B.P. 130/100. Pre-eclamptic symptoms
78	43	3	P	Surgical induction (bougies)	+	-	+	-	-	Esb. 5-12 parts. B.P. 190/120. Severe pre-eclamptic toxæmia
119	18	1	7	Undelivered -	+	-	-	-	-	Esbach, 12 parts. Pre-eclamptic symptoms
152	21	1	FT	Spontaneous -	+	-	+	-	-	Esbach, 3 parts. Pre-eclamptic symptoms
171	21	1	FT	Medical induction,	+	-	+	-	-	Esbach, 8 parts. Clear in 5 days
221	31	1	FT	Medical induction, spontaneous	+	-	+	-	-	Esbach, 1-8 parts. B.P. 140/100
230	22	2	FT	Spontaneous -	+	-	+	-	-	Alb. trace. B.P. 130/100
257	30	1	FT	Forceps -	+	-	+	-	-	Esbach, 2-12 parts. Pre-eclamptic symptoms
573	22	1	5	Undelivered -	+	-	-	-	-	B.P., 130/90. Esb., 1-2 parts

Reg. No.	Age.	Parity.	Stage of Pregnancy.	Mode of Delivery.	Result.					Remarks.	
					Mother.		Child.				
					L.	D.	L.	D.	S.B.		
574	30	2	FT	Spontaneous	-	+	-	+	-	-	Alb. trace. Mild case
601	40	6	FT	Spontaneous	-	+	-	+	-	-	Alb. trace. Mild case
657	—	9	FT	Manual	-	+	-	+	-	-	Esbach, 1 part. B.P. 160/120
668	31	7	FT	Spontaneous	-	+	-	+	-	-	Alb. trace. Mild case
671	21	2	6	Undelivered	-	+	-	-	-	-	Alb. trace. B.P. 120/90
796	30	2	8	Surgical induction (bougies)		+	-	+	-	-	Alb. trace. B.P. 150/100
805	38	7	8	Surgical induction (bougies)		+	-	-	-	+	Esb., 5 parts. B.P. 205/170. Chronic ne- phritis
889	38	9	8	Undelivered	-	+	-	-	-	-	Esbach, 2-3 parts. B.P. 200/150. Irregular dism.
933	34	4	8	Undelivered	-	+	-	-	-	-	Alb. trace. Irregular dism.
961	30	4	7	Surgical induction (bougies)		+	-	-	-	+	Esbach., 2-14 parts. B.P. 140/90
988	29	1	FT	Spontaneous	-	+	-	+	-	-	Esbach, 1-10 parts. B.P., 125/90
1009	31	1	8	Forceps	-	+	-	+	-	-	Esbach, 2-7 parts. B.P. 165/100
1021	26	4	8	Undelivered	-	+	-	-	-	-	Alb. trace. Mild case
1060	35	1	FT	Medical induction, manual		+	-	+	-	+	Esbach, 12 parts. B.P. 150/90
1077	37	6	8	Undelivered	-	+	-	-	-	-	Alb. trace. Mild case. B.P. 140/90
1173	29	2	FT	Surgical induction, spontaneous delivery (bougies)		+	-	+	-	-	Esb. 4-12 parts. B.P. 185/120. Pre-eclam. tox.
1256	28	1	FT	Forceps	-	+	-	+	-	-	B.P. 140/100. Esb. 1 part. Mild case
1262	29	2	—	Undelivered	-	+	-	+	-	-	B.P. 120/70. Esb. 1 part. Mild case
1274	34	6	8	Undelivered	-	+	-	-	-	-	B.P. 120/70. Alb. trace. Mild case
1320	30	4	6	Complete miscar- riage		+	-	-	-	-	Esb. 2-13 parts. B.P. 165/110
1343	22	1	FT	Medical induction, spontaneous		+	-	-	-	+	Esb., 1 part. B.P. 130/90. Pre-eclam. tox.
1345	38	5	FT	Surgical induction, spontaneous delivery (bougies)		+	-	+	-	-	Esb., 1 part. B.P. 230/140. Chronic ne- phritis
1374	24	2	7	Spontaneous	-	+	-	-	-	+	Esb. ½-5 parts. B.P. 130/90. Irregular dism.
1457	37	1	6	Undelivered	-	+	-	-	-	-	Esb. ½-10 parts. B.P. 200/160. Irregular dism.

Reg. No.	Age.	Parity.	Stage of Pregnancy.	Mode of Delivery.	Result.					Remarks.	
					Mother.		Child.				
					L.	D.	L.	D.	S.B.		
1458	26	2	FT	Spontaneous	-	+	-	+	-	-	Esb. 1 part. B.P. 120/65. Pre-eclam tox.
1471	28	3	7	Spontaneous	-	+	-	-	-	+	Esb. 1-7 parts. B.P. 140/80. Chronic nephritis
1472	23	2	FT	Spontaneous	-	+	-	+	-	-	Alb. trace. Mild case
1476	18	1	FT	Forceps	-	+	-	+	-	-	Esb. ½-2 parts. Pre-eclam. tox.
1515	33	1	FT	Spontaneous	-	+	-	+	-	-	Alb. trace. Mild case
1533	43	8	P	Spontaneous	-	+	-	-	-	+	Esb. 2-14 parts. B.P. 200/120. Chronic nephritis
1591	43	8	8	Undelivered	-	+	-	-	-	-	Alb. trace. B.P. 130/80. Mild case
1594	34	6	FT	Spontaneous	-	+	-	+	-	-	Esb. ½ part. Mild case
1642	27	2	7	Spontaneous	-	+	-	-	-	+	Esb. 1-7 parts. B.P. 190/110. Chronic nephritis
1654	26	2	FT	Spontaneous	-	+	-	++	+	-	Esb. ½ part. B.P. 150/90. Mild case
1664	42	5	FT	Spontaneous	-	+	-	+	-	-	Esb. 1 part. B.P. 180/90. Chronic nephritis
1674	30	3	FT	Manual extraction		+	-	+	-	-	Esb. 1 part. B.P. 120/80. Chronic nephritis
1710	25	1	7	Undelivered	-	+	-	-	-	-	Esb. 1-6 parts. B.P. 200/180
1741	31	1	8	Spontaneous	-	+	-	-	-	+	B.P. 170/105. Esb. 1-6 parts. Pre-eclam. tox.
1745	23	1	FT	Forceps	-	+	-	+	-	-	B.P. 140/90. Esb. ½ part. Pre-eclam. tox.
1760	21	1	7	Undelivered	-	+	-	-	-	-	Alb. trace. B.P., 120/70. Mild case
1789	15	1	FT	Spontaneous	-	+	-	+	-	-	Alb. trace. B.P., 110/70. Pre-eclam. symptoms
1800	40	5	FT	Spontaneous	-	+	-	+	-	-	Alb. trace. Marked œdema. Mild case
1804	43	9	FT	Spontaneous	-	+	-	+	-	-	Esb. ½-2 parts. B.P. 135/85
1810	20	1	8	Medical induction, spontaneous		+	-	+	-	-	Esb. 1-6 parts. B.P. 160/110
1835	19	1	FT	Spontaneous	-	+	-	+	-	-	Alb. trace. Mild case
1839	35	3	7	Spontaneous	-	+	-	-	-	+	Esb. 2-9 parts. B.P. 200/130. Conc. acc. hæm. Chronic nephritis
1888	36	2	FT	Spontaneous	-	+	-	+	-	-	Esb. 1 part. B.P. 120/80. Pre-eclam. tox.

Reg. No.	Age.	Parity.	Stage of Pregnancy.	Mode of Delivery.	Result.					Remarks.	
					Mother.		Child.				
					L.	D.	L.	D.	S.B.		
1910	35	3	FT	Spontaneous	-	+	-	++	-	-	Alb. trace. B.P. 140/90. Mild case
2193	27	4	—	Spontaneous	-	+	-	+	-	-	Esb. 2-6 parts. Chronic nephritis. B.P. 170/ 120
2241	29	1	FT	Manual	-	+	-	+	-	+	Esb. 2-9 parts. B.P. 160/130
2250	32	2	8	Spontaneous	-	+	-	+	-	-	B.P. 120/90. Alb. trace. Mild case
2276	27	3	8	Undelivered	-	+	-	-	-	-	Alb. trace. Mild case
2422	38	11	FT	Spontaneous	-	+	-	+	-	-	Alb. trace. B.P., 150/100. Mild case
2440	39	11	FT	Spontaneous	-	+	-	+	-	-	Esb. 2-8 parts. B.P. 190/100. Chronic ne- phritis
2461	24	1	FT	Spontaneous	-	+	-	+	-	-	Alb. trace. B.P., 170/120
2488	37	3	8	Undelivered	-	-	+	-	-	-	Mixed acc. hæm. Blood transfusion
2498	26	1	FT	Manual	-	+	-	+	-	-	B.P. 150/100. Esb. 2 parts. Pre-eclam. tox.
2509	28	1	FT	Spontaneous	-	+	-	+	-	-	Esb. 1-6 parts. B.P. 160/120
2512	32	4	FT	Spontaneous	-	+	-	+	-	-	Esb. 2-4 parts. B.P. 150/110
2565	28	1	FT	Spontaneous	-	+	-	+	-	-	Alb. trace. B.P. 140/90. Mild case
2566	27	4	—	Spontaneous	-	+	-	+	-	-	Alb. trace. B.P. 120/90. Mild case
2681	19	1	FT	Forceps	-	+	-	+	-	-	Esb. 1 part. B.P. 130/100
2686	22	1	FT	Spontaneous	-	+	-	+	-	-	Esb. 1-4 parts. B.P. 160/120
2708	23	1	FT	Spontaneous	-	+	-	+	-	-	Esb. 1 part. B.P., 110/70
2769	39	10	8½	Spontaneous	-	+	-	+	-	-	B.P. 200/120. Esb., 1-3 parts
2815	39	7	8	Undelivered	-	-	+	-	-	-	Mixed acc. hæm. Chronic Nephritis., Esb. 4½ parts
2823	26	4	7	Surgical induction, spontaneous (bougies)		+	-	+	+	-	B.P. 150/110. Chronic nephritis. Puer. mania.
2837	21	1	FT	Manual extraction (2)		+	-	++	-	-	Alb. trace. B.P. 120/90. Mild case
2845	28	2	FT	Spontaneous	-	+	-	+	-	-	Esb. 1 part. B.P., 110/70. Mild case
2862	26	1	FT	Medical induction, forceps		+	-	+	-	-	Esb. 2-7 parts. B.P., 160/90. Pre-eclam., tox.

Reg. No.	Age.	Parity.	Stage of Pregnancy.	Mode of Delivery.	Result.					Remarks.
					Mother.		Child.			
					L.	D.	L.	D.	S.B.	
2879	19	1	8	Spontaneous -	+	-	+	-	-	Esb. 2 parts. B.P. 125/80
2936	27	1	7	Surgical induction, (bougies), spontaneous	+	-	+	+	-	Esb. 4-10 parts. B.P. 230/160. Chronic nephritis
2964	35	4	8	Undelivered -	+	-	-	-	-	Trace alb. B.P., 160/90.
3011	41	4	FT	Spontaneous -	+	-	+	-	-	Esb. 2 parts. B.P. 150/100. Marked œdema
3033	32	1	FT	Manual extraction	+	-	+	-	-	Esb. 1-12 parts. B.P. 200/130. Pre-eclam. tox.
3043	25	4	FT	Spontaneous -	+	-	+	-	-	Alb. trace. B.P. 120/80. Mild case
3068	33	1	FT	Forceps - -	+	-	-	-	+	Esb. 2-10 parts. B.P. 180/100
3110	25	1	FT	Forceps - -	+	-	+	-	-	Alb. trace. B.P. 150/100
3122	31	1	8	Manual - -	+	-	-	-	+	4 fits. B.P. 190/130. Esb., 12 parts
3134	28	1	FT	Forceps (twins) -	+	-	+	-	+	Esb. 20 parts. B.P. 120/80. Comatose. 11 fits
3149	24	1	FT	Forceps - -	-	+	+	-	-	Trace alb. Death from pulmonary embolus.
3155	20	2	FT	Spontaneous -	+	-	+	-	-	Esb. 2 parts. B.P. 150/100. Chronic nephritis
3184	25	1	18	Spontaneous -	+	-	+	-	-	Alb. trace. Mild case. B.P. 110/70
3224	28	3	FT	B.B.A. - -	-	+	-	-	-	Esb. 16 parts. Chronic nephritis. Uræmia
3386	26	4	FT	Forceps - -	+	-	-	-	+	Esb. ½-2 parts. B.P. 180/130. F.F.O. Pre-eclam. tox.
3399	28	5	8	Spontaneous -	+	-	+	-	-	B.P., 130/90. Alb. trace. Mixed acc. hæm.
3496	18	1	FT	Spontaneous -	+	-	+	-	-	B.P. 170/130. Esb. 2-6 parts. Pre-eclam. tox.
3422	19	1	—	Forceps - -	+	-	+	-	-	Esb. 1 part. B.P. 180/120. Puer. sepsis. Transferred
3458	25	3	FT	Spontaneous -	+	-	+	-	-	Alb. trace. Mild case
3570	38	5	6	Surgical induction, Complete miscarriage	+	-	-	-	-	Esb. 15 parts. B.P. 200/100. Chronic nephritis
3593	35	1	8	Spontaneous -	+	-	+	+	-	Esb. 2 parts. B.P. 170/130
3608	—	4	5	Undelivered -	+	-	-	-	-	Alb. trace. B.P. 120/70. Mild case

Reg. No.	Age.	Parity.	Stage of Pregnancy.	Mode of Delivery.	Result.					Remarks.
					Mother.		Child.			
					L.	D.	L.	D.	S.B.	
3652	38	10	8	Spontaneous -	+	-	-	-	+	Esb. 1-6 parts. B.P. 210/100
3640	17	1	7½	Spontaneous -	+					Esb. 4-6 parts. B.P. 150/95
3652	38	10	FT	Spontaneous -	+	-	-	-	+	Esb. 1-6 parts. B.P. 210/100. Chronic nephritis
3673	29	2	6	Complete miscarriage	+	-	-	-	-	Esb. 2-8 parts. B.P. 190/150
3702	29	1	8	Spontaneous -	+	-	+	-	-	Esb. 7 parts. B.P. 160/120. Pre-eclam. tox.
3738	36	5	FT	Spontaneous -	+	-	-	-	+	Esb. ½ part. B.P. 130/80. Acc. hæm.
3741	21	1	6	Surgical induction Complete miscarriage	+	-	-	-	-	Esb. 12 parts. B.P. 160/110
3855	39	4	7	Spontaneous -	+	-	-	-	+	Esb. 2-12 parts. B.P. 180/110. Chronic nephritis. Acc. hæm.
3937	23	2	FT	Spontaneous -	+	-	+	-	-	Alb. trace. B.P. 140/70. Mild case
3951	40	4	6	Complete miscarriage	+	-				Esb. 4-12 parts. B.P. 180/30. Chronic nephritis
4009	42	4	6	Surgical induction Complete miscarriage	+	-	-	-	-	Esb. 2-8 parts. B.P. 240/145
4027	27	3	FT	Spontaneous -	+	+	+	-	-	Alb., trace. Mild case. Chronic nephritis
4045	24	13	4	Surgical induction Complete miscarriage	+	-	-	-	-	Chronic nephritis. Esb., 1 part. B.P. 230/170
4047	37	5	8	Spontaneous -	+	-	-	-	++	Esb. ½ part. B.P. 145/80 Conc. acc. hæm.
4079	34	8	8	Forceps - -	+	-	-	-	+	Alb. trace. Mild case
4089	19	1	8	Surgical induction, spontaneous	+	-	-	-	+	Esb. 2-10 parts. B.P. 130/100. Pre-eclam. tox.
4138	27	1	FT	Manual extraction	+	-	-	-	+	Esb. 2 parts. B.P. 170/100
4190	34	1	—	Forceps - -	+	-	-	-	+	Alb. trace. B.P. 130/80. Mild case
4239	23	6	7	Undelivered -	+	-	-	-	-	Esb. ½ part. B.P. 130/100
4309	28	3	4	Undelivered -	+	-	-	-	-	Chronic nephritis. Irregular dism.
4336	35	4	FT	Spontaneous -	+	-	+	-	-	Esb. 2-10 parts. B.P. 150/90.
4388	36	9	8	Spontaneous -	+	-	-	-	+	Esb. 13 parts. B.P. 150/80. Conc. acc. haem.

C.—Cases not under Ante-natal Supervision.

Reg. No.	Age.	Parity.	Stage of Pregnancy.	Mode of Delivery.	Result.					Remarks.	
					Mother.		Child.				
					L.	D.	L.	D.	S.B.		
316	39	8	FT	Spontaneous	-	+	-	+	-	-	Esbach 1 part.
642	38	7	8	Undelivered	-	+	-	-	-	-	Esbach 2 parts. Mild case.
751	29	1	8	Undelivered	-	+	-	-	-	-	Alb. trace. Pre-eclamptic symptoms.
756	21	3	8	Undelivered	-	+	-	-	-	-	Alb. trace. Mild case.
1283	34	7	7	Undelivered	-	+	-	-	-	-	Alb. trace. Mild case. B.P. 130/70.
1814	27	1	FT	Manual	-	+	-	-	-	+	Esb. 3 parts. B.P. 160/120. Pre-eclam. toxæmia.
2136	26	12	FT	Spontaneous	-	+	-	+	-	-	Alb. trace. Mild case.
2386	40	1	FT	Spontaneous	-	+	-	+	-	-	Esb. 3 parts. B.P. 150/100. Pre-eclam. tox.
2598	28	6	FT	Spontaneous	-	+	-	+	-	-	Esb. 13 parts. B.P. 180/90. Pre-eclam. tox.
2730	27	4	FT	Spontaneous	-	+	-	+	-	-	Esb. 1 part. B.P. 130/80.
3121	25	2	FT	Spontaneous	-	+	-	-	-	+	Esb. 4 parts. B.P. 170/90. Pre-eclamptic tox.
3562	36	1	5m	Miscarriage	-	+	-	-	-	+	Esb. 8 parts. B.P. 180/100. Pre-eclam. tox.
3573	22	1	FT	Spontaneous	-	+	-	+	-	-	Alb. trace. Mild case.
3841	40	6	FT	Spontaneous	-	+	-	+	-	-	Esb. 5 parts. B.P. 160/90. Pre-eclamp. tox.
4013	24	1	FT	Spontaneous	-	+	-	+	-	-	Marked œdema. Mild case.
4119	23	1	FT	Spontaneous	-	+	-	+	-	-	Alb. trace. Mild case.
4146	43	9	P	Spontaneous	-	+	-	+	-	-	Esb. 1 part. B.P. 170/120. Chronic nephritis.
4274	21	1	FT	Forceps	-	+	-	+	-	-	Esb. 1 part. B.P. 160/105.

ECLAMPSIA.

A.—Cases under Ante-natal Supervision.

Reg. No.	Age.	Parity.	Stage of Pregnancy.	Mode of Delivery.	Result.					Remarks.
					Mother.		Child.			
					L.	D.	L.	D.	S.B.	
1023	30	1	8	Undelivered -	-	+	-	-	-	32 fits. B.P. 170/100. Alb. ++.
1218	32	7	—	B.B.A. - -	+	-	-	-	-	2 fits. Esb. 8 parts. B.P. 155/115.
1492	16	1	FT	Forceps - -	+	-	+	-	-	3 fits. Esb. 3 parts. B.P. 118/82.
2639	19	2	8	Medical induction Spontaneous	+	-	-	-	+	3 fits. Esb. 13 parts. B.P. 130/90.
2656	22	1	FT	Forceps - -	+	-	+	-	-	9 fits. B.P. 180/90. Esb. parts.
2771	22	1	FT	Spontaneous -	+	-	+	-	-	4 fits. Esb. 8 parts. B.P. 165/120.
3485	41	5	8	Spontaneous -	+	-	+	-	-	4 fits. Esb. 13 parts. B.P. 210/110.

B.—Cases sent in as Emergencies.

326	25	1	FT	B.B.A. - -	+	-	-	-	-	Post. partum. B.P. 210/130. Esb. 7 parts. Comatose on admission.
459	34	1	7	Spontaneous -	+	-	-	-	+	Esb. 4-12 parts. Marked œdema.
504	22	1	FT	B.B.A. - -	+	-	-	-	-	Post. partum. B.P. 130/100. Esb. 2 parts. 3 fits.
602	27	2	P	Spontaneous -	+	-	+	-	-	Esb. 12 parts. 4 fits.
798	34	3	6	Miscarriage -	+	-	-	-	-	Esb. 4 parts. B.P. 195/120. Semi-comatose on admission. Good recovery.
1013	41	6	FT	B.B.A. - -	+	-	-	-	-	Post-partum eclampsia. B.P. 170/120. Esb. 3 parts.
1276	19	1	FT	Spontaneous -	+	-	+	-	-	3 fits. B.P. 155/95. Esb. 3 parts.
1306	32	1	FT	B.B.A. - -	+	-	-	-	-	3 fits. B.P. 175/110. Esb. 6 parts.
1308	30	1	7	Spontaneous -	+	-	-	-	+	1 fit. Esb. 15 parts. B.P. 170/120.
1331	42	3	7	Surg. ind. (bougies)	+	-	+	-	-	1 fit. Esb. 12-14 parts. B.P. 210/140.
1381	29	1	FT	Forceps - -	+	-	+	-	-	8 fits. Esb. 10 parts. B.P. 200/150.

Reg. No.	Age.	Parity.	Stage of Pregnancy.	Mode of Delivery.	Result.					Remarks.
					Mother.		Child.			
					L.	D.	L.	D.	S.B.	
1391	39	9	6	Surg. induction Complete mis- carriage	+	-	-	-	-	7 fts. Esb. 10 parts B.P. 205/150.
1466	33	1	FT	B.B.A. and Spontaneous	-	+	-	-	+	7 fts. B.P. 165/125. Esb. 8 parts. Intra- partum eclampsia.
1479	28	3	P	Spontaneous -	+	-	-	-	+	7 fts. B.P. 210/140. Comatose. Puerperal mania.
1572	36	1	8	Caes. section (post mortem)	-	+	-	-	++	11 fts. B.P. 120/80. Alb. ++.
1634	28	1	7½	Surg. induction (bougies)	+	-	+	+	-	2 fts. Esb. 8 parts. B.P. 215/120.
1692	30	1	6	Surg. induction (bougies) Miscarriage	+	-	+	-	-	3 fts. Alb. ++. B.P. 170/100.
1707	40	8	—	B.B.A. - -	+	-	-	-	-	10 fts. Alb. and blood ++. B.P. 145/100. Post-partum eclam- psia.
2184	20	1	8	Forceps - -	+	-	-	-	+	4 fts. Esb. 8 parts B.P. 140/90.
2205	26	1	FT	Forceps and Manual distrac- tion	-	+	-	-	++	10 fts. + Esb. 7 parts B.P. 160/180.
2244	28	3	FT	Forceps - -	+	-	+	-	-	1 fit. Esb. 8 parts. B.P. 160/100.
2258	22	1	FT	Forceps - -	+	-	+	-	-	6 fts. Esb. 4 parts. B.P. 140/80.
2430	25	2	FT	Spontaneous	+	-	-	-	+	11 fts. + Esb. 8 parts. B.P. 220/150.
2467	27	1	8	B.B.A. - -	-	+	+	-	-	2 fts. Esb. 12 parts. B. P. 150/100. Post- partum eclampsia.
2469	22	1	FT	Spontaneous -	+	-	+	-	-	5 fts. Esb. 12 parts. B.P. 170/118.
2476	20	1	—	B.B.A. - -	+	-	-	-	-	15 fts. Esb. 8 parts. B.P. 160/110.
2511	18	1	FT	Spontaneous -	-	+	+	-	-	8 fts. B.P. 150/110. Alb. ++ Cerebral hæmorrhage.
2634	17	1	FT	Spontaneous	+	-	-	-	+	1 fit. Esb. 12 parts. B.P. 120/70.
2664	36	3	FT	Undelivered -	-	+	-	-	-	6 fts. B.P. 165/95. Alb. ++.
2846	23	1	8	Forceps - -	+	-	+	-	-	5 fts. B.P. 150/110. Esb. 2 parts.
2932	40	7	FT	Spontaneous -	-	+	+	-	-	5 fts. B.P. 195/100. Marked jaundice. Alb. ++.

Reg. No.	Age.	Parity.	Stage of Pregnancy.	Mode of Delivery.	Result.					Remarks.
					Mother.		Child.			
					L.	D.	L.	D.	S.B.	
3122	31	1	8	Manual - -	+	-	-	-	+	4 fits. B.P. 190/130. Esb. 12 parts.
3134	28	1	FT	Forceps (twins)	+	-	+	-	+	Esb. 20 parts. B.P. 120/80. Comatose. 11 fits.
3135	21	1	74	Spontaneous -	+	-	+	-	-	8 fits. Esb. 15 parts. B. P. 140/110.
3161	33	1	FT	Forceps - -	+	-	-	-	+	11 fits. B.P. 230/150. Esb. 13 parts.
3219	35	7	FT	B.B.A. - -	+	-	-	-	-	5 fits. Esb. 3 parts. B.P. 170/110.
3325	19	1	FT	Forceps - -	+	-	+	-	-	1 fit. Esb. 12 parts. B.P. 190/120.
3407	35	1	8	Spontaneous -	+	-	-	-	+	7 fits. Esb. 24 parts. B.P. 190/125.
3482	18	1	8	Forceps - -	+	-	-	-	+	12 fits. Alb. + +. Maniacal. Transferred to Mental Hospital.
3664	34	3	6	Complete miscarriage	-	+	-	-	-	1 fit. B.P. 170/110. Pulmonary œdema.
3734	28	1	7	Spontaneous -	+	-	+	-	-	4 fits. Esb. 8 parts. B.P. 170/115.
3745	28	1	FT	Forceps - -	+	-	+	-	-	3 fits + Esb. 10 parts. B.P. 165/115.
3771	45	1	—		+	-				
3826	26	1	FT	Spontaneous -	-	+	-	-	+	11 fits. Alb. + +. B.P. 180/115.
3840	32	5	FT	B.B.A. - -	+	-	-	-	-	6 fits. Esb. 6 parts. B.P. 160/95. Postpartum eclampsia.
3959	33	3	P	Spontaneous	+	-	-	-	+	4 fits. Esb. 13 parts. B.P. 180/160.
4363	19	1	8	Forceps - -	+	-	+	-	-	4 fits. Esb. 10 parts. B.P. 190/120.
4453	39	1	8	Surg. induction (bougies)	+	-	-	-	-	4 fits. Esb. 20 parts. B.P. 200/100.
4469	27	1	8	Forceps - -	+	-	+	+	-	10 fits. Esb. 6 parts. B.P. 190/160.

C.—Cases not under Ante-natal Supervision.

1724	34	1	FT	Spontaneous -	+	-	+	-	-	1 fit. Esb. 1 part. B.P. 120/90.
2720	19	1	FT	Spontaneous -	+	-	+	-	-	2 fits. B.P. 164/120. Esb. 2 parts.
4125	23	1	FT	Spontaneous -	+	-	+	-	-	1 fit. Esb. 12 parts. B.P. 155/115.

HYPEREMESIS GRAVIDARUM AND LATE VOMITING.

A.—Cases under Ante-natal Supervision.

Reg. No.	Age.	Parity.	Stage of Pregnancy.	Mode of Delivery.	Result.					Remarks.
					Mother.		Child.			
					L.	D.	L.	D.	S.B.	
110	34	11	8	Surgical induction	+	-	-	-	+	Persistent late vomiting of pregnancy.
120	32	2	3	Undelivered -	+	-	-	-	-	Acetonuria. Recovered in 4 weeks.
137	26	4	6	Undelivered -	+	-	-	-	-	Late vomiting. Readmitted. Normal delivery at term.
175	21	1	4	Undelivered -	+	-	-	-	-	Mild case.
414	38	7	8	Undelivered -	+	-	-	-	-	Late vomiting. Re-admitted at term. Normal delivery.
786	28	5	4	Undelivered -	+	-	-	-	-	Mild case. Acetonuria.
831	33	10	3	Undelivered -	+	-	-	-	-	Mild case. Acetonuria.
1062	34	8	3	Undelivered -	+	-	-	-	-	Mild case. Acetonuria.
1160	24	4	7	Spontaneous -	+	-	-	-	+	Late vomiting of pregnancy.
1576	42	8	2	Miscarriage -	+	-	-	-	-	Persistent vomiting. Good recovery.
1822	25	1	2	Undelivered -	+	-	-	-	-	Mild case. Good recovery.
1919	37	4	3	Undelivered -	+	-	-	-	-	Mild case. Good recovery.
2009	28	2	3	Undelivered -	+	-	-	-	-	R.G.U. Replaced. Vomiting stopped.
2057	25	3	3½	Undelivered -	+	-	-	-	-	Neurotic vomiting. Good recovery.
2583	24	1	2	Undelivered -	+	-	-	-	-	Mild case. Good recovery.
2887	30	5	2½	Undelivered -	+	-	-	-	-	Marked constipation. Severe vomiting. Acetonuria.
3008	21	1	8	Undelivered -	+	-	-	-	-	Late vomiting of pregnancy. Good recovery.
3094	24	1	3	Undelivered -	+	-	-	-	-	Neurotic vomiting. Good recovery.
3232	30	1	3	Undelivered -	+	-	-	-	-	Neurotic vomiting. Rather emaciated. Good recovery.
3389	26	2	4	Undelivered -	+	-	-	-	-	Emaciated. Marked Acetonuria. Achlorhydria.
3440	29	2	2	Undelivered -	+	-	-	-	-	Marked neurotic vomiting. 8 weeks' treatment. Good recovery.

Reg. No.	Age.	Parity.	Stage of Pregnancy.	Mode of Delivery.	Result.					Remarks.	
					Mother.		Child.				
					L.	D.	L.	D.	S.B.		
3468	22	2	6	Undelivered	-	+	-	-	-	-	Neurotic vomiting. Mild case.
3687	22	3	5	Undelivered	-	+	-	-	-	-	Constipation severe. Good recovery.
3761	29	3	8½	Undelivered	-	+	-	-	-	-	Late vomiting. Good recovery.
3892	22	1	3	Undelivered	-	+	-	-	-	-	Neurotic vomiting Good recovery.
3942	29	2	2½	Undelivered	-	+	-	-	-	-	Persistent vomiting. Acetonuria. Toxic. Good recovery after 9 weeks.
4086	24	1	3	Undelivered	-	+	-	-	-	-	Emaciated. Acetonuria. Good recovery.
4087	30	3	3	Undelivered	-	+	-	-	-	-	Persistent neurotic vomiting. Good recovery.
4091	27	4	6	Undelivered	-	+	-	-	-	-	Neurotic vomiting.
4128	20	1	3	Undelivered	-	+	-	-	-	-	Neurotic vomiting. Good recovery.
4277	34	11	4	Undelivered	-	+	-	-	-	-	Neurotic vomiting.
4373	24	1	3	Undelivered	-	+	-	-	-	-	Acetonuria. Dehy- dration. Good re- covery.

B.—Cases sent in as Emergencies

20	27	3	3	Undelivered	-	+	-	-	-	Acetonuria. Moderately severe case. Recovered in 3 weeks.
84	19	1	6	Complete miscarriage		+	-	-	-	Moderately severe case. 5 weeks' treatment.
158	25	1	3	Induction of abortion		-	+	-	-	Dehydrated. Jaundiced. Pulse 140 on admission.
256	23	6	6	Undelivered	-	+	-	-	-	3 months' sickness. Marked constipation.
298	23	2	3	Undelivered	-	+	-	-	-	Moderately severe case. Acetonuria. Slight jaundice.
350	25	3	4	Undelivered	-	+	-	-	-	Mild case.
401	24	3	6	Undelivered	-	+	-	-	-	Marked constipation.
450	38	5	4	Undelivered	-	+	-			Acetonuria. Pulse 120 on admission. Recovered in 2 weeks.
553	19	1	3	Undelivered	-	+	-	-	-	Mild case.
587	31	2	2	Induction of abortion (lamenarial tents)		+	-	-	-	Abortion induced after 6 weeks medical treatment.
901	33	5	8	Undelivered	-	+	-	-	-	Recovered in 2 weeks.
943	19	1	4	Undelivered	-	+	-	-	-	Mild case.

Reg. No.	Age.	Parity.	Stage of Pregnancy.	Mode of Delivery.	Result.					Remarks.
					Mother.		Child.			
					L.	D.	L.	D.	S.B.	
958	32	4	3	Undelivered -	+	-	-	-	-	Dehydrated. Pulse 110. Acetonuria. 4 weeks' treatment.
1039	35	3	3	Complete abortion	-	+	-	-	-	Severe toxæmia. Died 6 weeks later.
1046	37	3	FT	Surgical induction	+	-	-	-	+	Late vomiting. Epigastric pain. Carpopedal spasm.
1072	24	2	3	Undelivered -	+	-	-	-	-	Neurotic type.
1110	35	8	3	Undelivered -	+	-	-	-	-	Retroverted gravid uterus. Corrected.
1163	27	5	4	Undelivered -	+	-	-	-	-	Neurotic vomiting. Mild case.
1201	18	1	6	Undelivered -	+	-	-	-	-	Severe vomiting. Slight jaundice. Acetonuria. Recovered in 5 weeks.
1239	24	1	2	Undelivered -	+	-	-	-	-	Marked constipation. Neurotic vomiting. Delivery of twins (forceps) at term.
1251	38	8	3	Undelivered -	+	-	-	-	-	Neurotic vomiting.
1267	21	1	7	Surgical induction	+	-	-	-	+	Emaciated. Toxic. 12 weeks in hospital.
1355	32	3	7½	Spontaneous -	+	-	-	-	+	Late vomiting. Persistent until delivery.
1455	38	1	5	Surgical induction (bougies)	-	+	-	-	-	Toxic on admission. Pulse 130. Jaundice. 3½ weeks in hospital.
1522	28	2	3	Miscarriage Undelivered -	+	-	-	-	-	Neurotic vomiting. Readmitted and delivered (forceps) at term.
1597	30	4	7	Spontaneous -	+	-	+	-	-	Late vomiting. Good recovery.
1619	21	1	3	Undelivered -	+	-	-	-	-	Severe vomiting. Dehydrated. Acetonuria. Intravenous glucose, etc. Readmitted at term. Normal delivery.
1693	34	7	3	Undelivered -	+	-	-	-	-	Neurotic vomiting.
1790	34	2	4	Induction of abortion	+	-	-	-	-	Persistent vomiting. Slight jaundice. 7 weeks in hospital.
1988	26	1	2	Undelivered -	+	-	-	-	-	Mild case.
1989	25	1	2	Undelivered -	+	-	-	-	-	Constipated. Mild case.
2177	23	1	2	Undelivered -	+	-	-	-	-	Hysterical patient. Good recovery.
2251	25	2	FT	Med. induction Spontaneous	+	-	+	-	-	Late vomiting.

Reg. No.	Age.	Parity.	Stage of Pregnancy.	Mode of Delivery.	Result.					Remarks.
					Mother.		Child.			
					L.	D.	L.	D.	S.B.	
2278	19	1	8	Surgical induction	+	-	+	-	-	Persistent vomiting.
				Spontaneous						Slight jaundice. Toxic.
2338	32	3	3	Undelivered	-	+	-	-	-	Mild case.
2356	25	2	4	Undelivered	-	+	-	-	-	Neurotic vomiting.
										Good recovery.
2357	21	1	2	Undelivered	-	+	-	-	-	Dehydrated. Acetonuria.
										5 weeks treatment. Good recovery.
2371	27	3	3	Undelivered	-	+	-	-	-	Neurotic vomiting.
2486	21	1	5	Undelivered	-	+	-	-	-	Severe vomiting.
										Dehydration.
										Acetonuria.
2673	39	4	3	Undelivered	-	+	-	-	-	Emaciation. Acetonuria.
										Severe vomiting
										Recovered in 4 weeks.
2674	22	1	6	Undelivered	-	+	-	-	-	Late vomiting.
										Acetonuria.
2697	36	6	7	Undelivered	-	+	-	-	-	Late vomiting.
										Glycosuria.
2731	25	2	3	Undelivered	-	+	-	-	-	Neurotic vomiting.
										Good recovery.
2746	26	3	36 wks	Surgical induction	+	-	+	-	-	Toxic vomiting.
				Spontaneous						Jaundice and visual disturbance.
2752	24	2	2	Undelivered	-	+	-	-	-	Continuous vomiting.
										Good recovery.
2763	29	2	8	Spontaneous	-	+	-	+	-	Late vomiting until delivery.
2792	26	2	3	Undelivered	-	+	-	-	-	Marked neurotic vomiting.
										Good recovery.
2825	29	1	3	Uterus evacuated (hydat. mole)	+	-	-	-	-	Severe vomiting.
2928	23	4	2	Undelivered	-	+	-	-	-	Mild case.
3151	24	1	3	Abdominal hysterotomy	-	-	+	-	-	Emaciated and cachectic. Marked toxæmia.
3153	22	2	3	Undelivered	-	+	-	-	-	Mild case.
3310	15	1	4	Abdominal hysterotomy	+	-	-	-	-	Toxic vomiting.
3379	29	1	6 wks	Undelivered	-	+	-	-	-	Mild case.
3379	29	1	3	Undelivered	-	+	-	-	-	Mild case.
3381	31	7	3	Undelivered	-	+	-	-	-	Mild case.
3382	21	1	3	Undelivered	-	+	-	-	-	Toxic vomiting.
										Irregular dismissal.
3450	23	2	4	Undelivered	-	+	-	-	-	Severe constipation.
										Good recovery.
3454	31	2	4	Abdominal hysterotomy	-	-	+	-	-	Severe toxæmia.
										Jaundice. Emaciation.
										Persistent vomiting.
3501	29	3	3	Undelivered	-	+	-	-	-	Neurotic vomiting.
										Dehydration. Slight jaundice.

Reg. No.	Age.	Parity.	Stage of Pregnancy.	Mode of Delivery.	Result.					Remarks.	
					Mother.		Child.				
					L.	D.	L.	D.	S.B.		
3572	36	2	2	Undelivered	-	+	-	-	-	-	Mild case.
3653	22	1	3	Undelivered	-	+	-	-	-	-	Mild case.
3856	39	7	3	Undelivered	-	+	-	-	-	-	Dehydrated. Excessive sickness.
3949	21	1	3	Undelivered	-	+	-	-	-	-	Mild case. Acetonuria.
3963	25	1	3	Undelivered	-	+	-	-	-	-	Marked neurotic vomiting.
4085	25	3	3	Induction of abortion.		+	-	-	-	-	Persistent vomiting.
4120	35	10	3	Undelivered	-	+	-	-	-	-	Emaciation. Jaundice.
4145	21	1	3	Undelivered	-	+	-	+	-	-	Marked neurotic vomiting.
											Severe vomiting. Re-admitted for 5 weeks at term. Normal delivery.
4179	28	5	3	Undelivered	-	+	-	-	-	-	Mild case.
4203	22	1	3	Undelivered	-	+	-	-	-	-	Emaciation and dehydration. Recovered in 6 weeks.
4223	28	5	3	Undelivered	-	+	-	-	-	-	Mild case.
4240	38	3	3	Undelivered	-	+	-	-	-	-	Mild case.
4248	20	1	7	Undelivered	-	+	-	-	-	-	Mild case. Late vomiting.
4334	22	3	4	Undelivered	-	-	-	-	-	-	Mild case. Neurotic vomiting.
4376	23	2	2	Undelivered	-	+	-	-	-	-	Persistent vomiting.
											Recovered in 5 weeks.
4413	32	3	3	Undelivered	-	+	-	-	-	-	Neurotic vomiting.

C.—Cases not under Ante-natal Supervision.

987	28	5	8	Surgical induction	-	+	-	-	+	Late vomiting. No ante-natal care. Seriously ill on admission.
2031	22	2	8	Spontaneous	-	+	-	-	+	Late vomiting.
3889	27	7	6	Undelivered	-	+	-	-	-	Mild case. Late vomiting.

PYELITIS.

*Table—Cases of Pyelitis admitted during 1930.**A—Cases under Ante-natal Supervision.*

During the year 1930 there were 132 cases of pyelitis admitted. Of these, 51 were in Category A, 65 in Category B, and 16 in Category C.

Reg. No.	Age.	Parity.	Stage of Pregnancy.	Delivery.	Result.					Miscarriage.	Remarks.	
					Mother.		Child.					
					A.	D.	A.	S.B.	D.			
176	18	2	FT	Spontaneous	-	+	-	+	-	-	-	4 weeks treatment. Good recovery
211	21	1	6	Undelivered	-	+	-	-	-	-	-	15 days treatment
476	27	6	FT	Spontaneous	-	+	-	+	-	-	-	14 days treatment Good recovery
515	28	4	FT	Spontaneous	-	+	-	+	-	-	-	Severe P.P.H.
753	31	7	8	Undelivered	-	+	-	-	-	-	-	
900	40	4	6	Miscarriage	-	+	-	-	-	-	+	9 weeks treatment
932	28	1	9	Forceps	-	+	-	+	-	-	-	
1063	31	3	8	Undelivered	-	+	-	-	-	-	-	B.P. 156/90
1107	28	1	8	Spontaneous		+	-	++	-	-	-	Prem. labour
				(twins)								
1713	26	3	8	Spontaneous		+	-	++	-	-	-	Prem. labour
				(twins)								
1740	19	1	8	Undelivered	-	+	-	-	-	-	-	
1980	19	1	6	Undelivered	-	+	-	-	-	-	-	Mild case
2037	30	1	8	Undelivered	-	+	-	-	-	-	-	Severe case
2042	18	1	6	Undelivered	-	+	-	-	-	-	-	Mild case
2044	22	3	6	Undelivered	-	+	-	-	-	-	-	Mild case
2045	30	7	7	Undelivered	-	+	-	-	-	-	-	Mild case
2046	27	7	3	Undelivered	-	+	-	-	-	-	-	Mild case
2093	40	10	FT	Spontaneous	-	+	-	+	-	-	-	
2141	23	1	FT	Forceps	-	+	-	+	-	-	-	Ureteral catheters, etc. 18 weeks treatment
2256	32	9	8	Undelivered	-	+	-	-	-	-	-	Mild case
2308	21	2	5	Undelivered	-	+	-	-	-	-	-	Mild case
2423	18	1	3	Undelivered	-	+	-	-	-	-	-	4 weeks treatment. Ureteral catheters, etc.
2497	20	1	FT	Spontaneous	-	+	-	+	-	-	-	Mild case
2568	34	7	FT	Spontaneous	-	+	-	+	-	-	-	12 weeks treatment. Good recovery.
2831	24	1	6	Undelivered	-	+	-	-	-	-	-	Mild case
2839	24	1	FT	Spontaneous	-	+	-	+	-	-	-	6 weeks treatment Good recovery
2863	25	3	FT	Spontaneous	-	+	-	+	-	-	-	3 weeks treatment. Good recovery
2886	24	1	FT	Spontaneous	-	+	-	+	-	-	-	
2910	27	3	8	Undelivered	-	+	-	-	-	-	-	4 weeks treatment. Good recovery
3076	26	4	6	Undelivered	-	+	-	-	-	-	-	Mild case
3120	34	6	FT	Undelivered	-	+	-	-	-	-	-	Irreg. dism.
3267	34	5	7	Undelivered	-	+	-	-	-	-	-	3 weeks treatment. Good recovery
3329	28	3	3	Undelivered	-	+	-	-	-	-	-	Mild case
3331	28	4	5	Undelivered	-	+	-	-	-	-	-	4 weeks treatment

Reg. No.	Age.	Parity.	Stage of Pregnancy.	Delivery.	Result.					Miscarriage.	Remarks.	
					Mother.		Child.					
					A.	D.	A.	S.B.	D.			
3397	38	6	6	Undelivered	-	+	-	-	-	-	-	3 weeks treatment. Good recovery
3415	27	3	6	Miscarriage	-	-	+	-	-	-	+	Ureteral catheters. P.M. Pyelonephrosis
3500	33	7	6	Miscarriage	-	+	-	-	-	-	-	10 days treatment before miscarriage
3520	19	1	FT	Spontaneous	-	+	-	-	-	-	-	Mild case
3557	35	9	6	Undelivered	-	+	-	-	-	-	-	Mild case
3581	23	1	4	Undelivered	-	+	-	-	-	-	-	4 weeks treatment. Good recovery
3595	25	3	5	Undelivered	-	+	-	-	-	-	-	
3641	19	1	5	Undelivered	-	+	-	-	-	-	-	7 weeks treatment. Good recovery
3698	18	1	6	Undelivered	-	+	-	-	-	-	-	Mild case
3907	25	1	FT	Spontaneous	-	+	-	+	-	-	-	Mild case
3964	26	4	6	Undelivered	-	+	-	-	-	-	-	Mild case
4182	19	1	7	Undelivered	-	+	-	-	-	-	-	
4255	20	1	7	Spontaneous	-	+	-	+	-	-	-	Severe case. Ureteral catheters, etc. 5 weeks treatment. Good recovery
4275	22	1	7	Undelivered	-	+	-	-	-	-	-	Mild case
4283	34	8	6	Undelivered	-	+	-	-	-	-	-	15 days treatment. Good recovery
4403	32	4	5	Undelivered	-	+	-	-	-	-	-	Mild case
4472	25	1	FT	Medical induction	-	+	-	+	-	-	-	Good recovery
				Spontaneous								

B.—Cases sent in as Emergencies.

28	32	2	8	Undelivered	-	+	-	-	-	-	-	18 days treatment
31	35	9	5	Undelivered	-	+	-	-	-	-	-	12 days in hospital. Good recovery
138	20	1	FT	Spontaneous	-	+	-	+	-	-	-	5 weeks treatment. P.P.H.
241	23	1	4	Undelivered	-	+	-	-	-	-	-	17 days treatment. Good recovery
277	21	1	4	Undelivered	-	+	-	-	-	-	-	Fortnights treatment. Good recovery
327	28	3	5	Undelivered	-	+	-	-	-	-	-	
440	25	3	8	Undelivered	-	+	-	-	-	-	-	Fortnights treatment. Good recovery
475	18	1	5	Undelivered	-	+	-	-	-	-	-	3 weeks treatment. Good recovery
478	20	2	4	Undelivered	-	+	-	-	-	-	-	14 days in hospital. Good recovery
517	35	6	8	Manual	-	+	-	+	-	-	-	Prem. labour
551	27	2	6	Undelivered	-	+	-	-	-	-	-	4 weeks treatment. Ureteral catheters, etc. Good recovery

Reg. No.	Age.	Parity.	Stage of Pregnancy.	Delivery.	Result.					Miscarriage.	Remarks.	
					Mother.		Child.					
					A.	D.	A.	S.B.	D.			
559	28	3	5	Undelivered	-	+	-	-	-	-		
572	31	5	7	Undelivered	-	+	-	-	-	-		
573	22	1	5	Undelivered	-	+	-	-	-	-	Irreg. dism.	
652	28	1	8	Spontaneous	-	+	-	+	-	-	Prem. labour	
696	23	1	6	Spontaneous	-	+	-	-	-	+	Prem. labour. 8 weeks treatment. Alkalies and catheters	
771	34	3	3	Induction of abortion		+	-	-	-	-	+	Marked toxæmia. 7 weeks treatment
976	40	2	FT	Undelivered	-	+	-	-	-	-	-	
1427	30	3	8	Spontaneous	-	+	-	+	-	-	-	Prem. labour
1444	19	1	7	Undelivered	-	+	-	-	-	-	-	3 weeks treatment. Good recovery
1488	23	1	6	Undelivered	-	+	-	-	-	-	-	3 weeks treatment
1507	35	13	7	Manual	-	+	-	+	-	-	-	3 weeks treatment
1582	37	8	7	Undelivered	-	+	-	-	-	-	-	Mild case
1726	21	3	FT	Spontaneous	-	+	-	+	-	-	-	7 weeks treatment. Good recovery
1729	23	1	7	Undelivered	-	+	-	-	-	-	-	Mild case
1732	39	8	8	Spontaneous	-	+	-	+	-	-	-	6 weeks treatment
1769	24	2	8	Undelivered	-	+	-	-	-	-	-	Sent in as appendicitis
1819	25	1	5	Spontaneous	-	+	-	-	-	-	+	Prem. labour. Severe case
1891	24	1	6	Undelivered	-	+	-	-	-	-	-	3 weeks treatment. Good recovery
1948	18	1	8	Undelivered	-	+	-	-	-	-	-	Mild case
2033	32	6	4	Undelivered	-	+	-	-	-	-	-	Mild case
2054	31	1	6	Miscarriage	-	+	-	-	-	-	-	3 weeks treatment
2076	25	1	4	Undelivered	-	+	-	-	-	-	-	Cystoscopy, etc.
2092	23	1	8	Undelivered	-	+	-	-	-	-	-	5 weeks treatment
2192	21	3	3	Undelivered	-	+	-	-	-	-	-	Fortnights treatment. Good recovery
2253	23	1	5	Undelivered	-	+	-	-	-	-	-	
2255	21	1	5	Undelivered	-	+	-	-	-	-	-	Mild case
2320	22	1	FT	Spontaneous	-	+	-	-	-	-	+	3 weeks treatment
2567	28	5	FT	Medical induction		+	-	+	-	-	-	6 weeks treatment
				Spontaneous delivery								Good recovery
2601	25	7	6	Miscarriage	-	+	-	-	-	-	+	Severe case. 8 weeks treatment
2707	23	1	8	Spontaneous	-	+	-	+	-	-	-	Severe case. Uret- eral catheters, etc.
2865	32	3	8	Undelivered	-	+	-	-	-	-	-	Mild case
3051	38	13	8	Undelivered	-	+	-	-	-	-	-	15 days treatment. Good recovery
3054	33	3	8	Undelivered	-	+	-	-	-	-	-	
3127	41	5	9	Undelivered	-	+	-	-	-	-	-	Mild case
3129	32	2	4	Undelivered	-	+	-	-	-	-	-	Mild case
3152	23	2	5	Undelivered	-	+	-	-	-	-	-	3 weeks treatment. Good recovery

Reg. No.	Age.	Parity.	Stage of Pregnancy.	Delivery.	Result.					Miscariage.	Remarks.	
					Mother.		Child.					
					A.	D.	A.	S.B.	D.			
3202	24	1	8	Undelivered	-	+	-	-	-	-	3 weeks treatment	
3231	24	1	7	Undelivered	-	+	-	-	-	-		
3456	23	4	FT	Forceps	-	+	-	+	-	-	Mild case	
3502	33	8	7	Undelivered	-	+	-	-	-	-	Irreg. dism.	
3506	27	1	5	Undelivered	-	+	-	-	-	-	3 weeks treatment	
3594	29	1	5	Undelivered	-	+	-	-	-	-	Mild case	
3725	25	4	3	Undelivered	-	+	-	-	-	-	Mild case	
3740	24	5	6	Undelivered	-	+	-	-	-	-	Mild case	
3875	28	3	8	Undelivered	-	+	-	-	-	-	15 days treatment.	
											Good recovery	
3979	30	5	8	Undelivered	-	+	-	-	-	-	Cystoscopy, etc. 2 weeks treatment.	
											Good recovery	
4093	33	4	FT	Spontaneous	-	+	-	+	-	-	Good recovery after 4 weeks treatment	
4162	35	9	6	Undelivered	-	+	-	-	-	-	Mild case	
4321	23	1	7	Undelivered	-	+	-	-	-	-	Severe case. Ureteral catheters, etc. 3 weeks treatment	
4351	21	3	8	Undelivered	-	+	-	-	-	-	Mild case	
4369	31	10	4	Undelivered	-	+	-	-	-	-	Mild case	
4391	43	13	6	Undelivered	-	+	-	-	-	-	Mild case	
4392	20	2	6	Undelivered	-	+	-	-	-	-	Ureteral catheters inserted. Good recovery	
4429	21	1	5	Undelivered	-	+	-	-	-	-	Mild case	
C.—Cases not under Ante-natal Supervision.												
43	34	6	7	Undelivered	-	+	-	-	-	-	10 days treatment	
139	26	2	FT	Spontaneous	-	+	-	+	-	-	14 days treatment	
279	18	1	8	Undelivered	-	+	-	-	-	-	Fortnights treatment. Good recovery	
471	22	1	8	Spontaneous	-	+	-	+	-	-	Pyuria and hæmaturia. Cystitis	
485	19	1	5	Surgical induction (bougies)		+	-	-	-	+	Severe toxæmia	
491	30	3	FT	Spontaneous	-	+	-	+	-	-	7 weeks treatment	
											Alkalies and ureteral catheters	
1035	39	9	8	Undelivered	-	+	-	-	-	-		
1042	31	5	7	Undelivered	-	+	-	-	-	-	Mild case	
1539	20	1	6	Undelivered	-	+	-	-	-	-	Mild case	
1860	22	1	7	Undelivered	-	+	-	-	-	-	Mild case	
2106	23	2	FT	Spontaneous	-	+	-	+	-	-	Good recovery	
2462	34	6	FT	Spontaneous	-	+	-	+	-	-	Cystoscopy, etc.	
											Severe case	
2551	21	1	5	Undelivered	-	+	-	-	-	-	Mild case	
2576	38	8	FT	Spontaneous	-	+	-	+	-	-		
3689	20	1	5	Undelivered	-	+	-	-	-	-	3 weeks treatment.	
											Good recovery	
3739	20	1	6	Undelivered	-	+	-	-	-	-	Mild case	

HEART DISEASE.

Table—Cases of Heart Disease admitted during 1930.

One hundred and thirteen cases of heart disease were admitted during the year 1930. Of these, 71 were under ante-natal supervision, 38 were sent in as emergencies, and 4 applied for admission without having been under previous supervision. There were 9 maternal deaths, 5 of which were in Category A and 4 in Category B.

Note.—The American Heart Association Functional Classification has been adopted in the following table. Class 1 denotes patients with organic heart disease who show no symptoms of discomfort. Class 2*a* embraces those cases which have fairly good compensation. In Class 2*b* there is some degree of decompensation, and in Class 3 the patients are definitely decompensated.

A.—Cases under Ante-natal Supervision.

Reg. No.	Age.	Parity.	Class.	Delivery.		State of Pregnancy on Admission.	Result.		Remarks.
							M.	Ch.	
165	39	3	2 <i>a</i>	Undelivered	-	5	A	—	4 weeks treatment. Much improved
167	21	1	1	Spontaneous	-	FT	A	A	Mitral incompetence. Mild case
174	25	3	2 <i>b</i>	Spontaneous	-	8	A	A	Mitral stenosis. Orthopnoea
190	35	12	1	Undelivered	-	6	A	—	3 weeks treatment. Much improved. Premature delivery (forceps) at term
192	35	7	1	Spontaneous (premature)		6	A	A	9 weeks treatment in all. Marked secondary anæmia
210	26	2	1	Undelivered	-	7	A	—	Mild case
335	44	13	2 <i>a</i>	Spontaneous	-	7	A	A	3 weeks ante-natal hospital treatment. Much improved. Readmitted for delivery
337	29	6	1	Undelivered	-	8	—	—	Well compensated mitral stenosis
393	26	5	2 <i>b</i>	Cæsarean section	-	5	A	A	Mitral incompetence. Orthopnoea, etc. 6 weeks ante-natal treatment. Sterilised
483	38	8	2 <i>a</i>	Cæsarean section	-	6	A	A	4 weeks ante-natal treatment. Readmitted. Cæsarean section. 38th week. Sterilisation.
518	23	2	2 <i>a</i>	Spontaneous	-	FT	A	A	Easy labour. Good recovery
641	27	3	2 <i>a</i>	Undelivered	-	6	A	—	Well compensated. Mitral stenosis

Reg. No.	Age.	Parity.	Class.	Delivery.	State of Pregnancy on Admission.	Result.		Remarks.
						M.	Ch.	
721	25	1	1	Spontaneous -	FT	A	A	Easy labour. Good recovery
795	18	1	1	Spontaneous -	FT	A	A	Easy labour. Good recovery
832	29	1	2b	Cæsarean section -	3	A	A	5 months ante-natal treatment. Sterilised
840	24	4	1	Spontaneous -	FT	A	A	Easy labour. Good recovery
939	33	5	2a	Forceps -	4	A	A	2 weeks ante-natal care. Delivered at term
1001	42	8	2a	Spontaneous -	FT	A	A	Mitral stenosis. Well compensated. Easy labour
1323	37	9	2b	Undelivered -	8	A	—	4 weeks ante-natal treatment. Much improved
1362	25	3	1	Undelivered -	5	A	—	Mild case
1470	26	3	1	Spontaneous -	FT	A	A	Marked secondary anæmia
1580	26	1	2a	Cæsarean section -	8	A	A	4 weeks ante-natal treatment. Not sterilised
1631	35	12	2a	Undelivered -	7	A	—	Mitral stenosis. Much improved
1715	24	4	1	Spontaneous -	7	A	A	2 weeks ante-natal treatment. Well compensated. Mitral stenosis
1911	?	4	2b	Cæsarean section -	8	A	A	6 weeks ante-natal treatment. Sterilisation
2041	27	1	1	Spontaneous -	6	A	A	3 weeks ante-natal treatment. Neurotic
2097	36	5	2b	Cæsarean section -	6	A	A	6 weeks ante-natal treatment. Complicated by chorea of pregnancy
2100	28	1	2b	Undelivered -	6	A	—	Complicated by chorea of pregnancy. Irreg. dismissal
2158	40	13	2a	Cæsarean section -	FT	A	A	Well compensated. Mitral stenosis. Sterilised
2166	34	9	2a	Spontaneous -	8	A	A	W.R. + Easy labour
2335	32	7	2b	Spontaneous -	7	A	A	Precipitate labour. 9 weeks ante-natal treatment. Improved considerably
2337	37	8	1	Manual (premature)	7	A	D	1 weeks ante-natal care. Delivered at term
2372	23	3	1	Spontaneous -	8	A	A	1 weeks ante-natal treatment. Easy labour at term
2481	41	10	2a	Spontaneous -	8	A	A	Well compensated. Mitral stenosis. Easy labour
2627	37	7	2b	Termination of pregnancy	3	A	—	Sea tangle tents. Induced abortion. Severe case
2628	29	4	2a	Undelivered -	8	—	—	4 weeks ante-natal treatment. Easy labour at home
2703	39	8	2b	Forceps -	7	A	A	11 weeks treatment. Good recovery
2733	27	4	2b	Spontaneous -	—	D	A	Puerperal ulcerative endocarditis. Chronic mitral stenosis
2741	19	1	1	Spontaneous (premature)	7	A	A	Mild case
2853	33	5	2a	Cæsarean section -	8½	A	A	Sterilised. Mitral stenosis

Reg. No.	Age.	Parity.	Class.	Delivery.	Stage of Pregnancy on Admission.	Result.		Remarks.
						M.	Ch.	
2899	36	7	2a	Cæsarean section -	8½	A	A	Well compensated. Mitral stenosis. Sterilised
2935	28	2	2a	Cæsarean section -	7½	A	A	Well compensated. Mitral incompetence. Sterilised
3009	33	6	2b	Cæsarean section -	8½	D	A	Sterilised. Mitral stenosis. Some loss of reserve
3093	33	7	2b	Spontaneous -	8½	A	A	Easy labour
3321	27	3	1	Spontaneous -	FT	A	A	Mild case
3330	20	3	1	Spontaneous -	7	A	A	2 weeks ante-natal treatment. Delivered at term
3441	27	1	2b	Cæsarean section -	3	D	A	Sterilisation. Mitral stenosis. Some loss of reserve. 5 Months treatment
3582	24	1	1	Undelivered -	7	A	—	Mild case
3583	21	1	3	Forceps -	8	A	A	Decompensated. Mitral stenosis. Marked improvement on treatment
3686	38	1	2a	Cæsarean section -	8½	A	A	Sterilised. Well compensated. Mitral stenosis
3760	23	1	2a	Spontaneous -	FT	A	A	Well compensated. Mitral stenosis
3796	40	10	2b	Spontaneous -	8	A	A	Fair amount of loss of reserve
3903	17	1	1	Cæsarean section -	8½	A	A	Operation on account of contracted pelvis
3908	41	9	2b	Undelivered -	3	A	—	Much improved after 3 weeks rest
3912	21	1	2b	Spontaneous (premature)	7½	A	A	Complicated by severe albuminuria
3989	27	1	2b	Termination of pregnancy	3	A	—	Loss of compensation. Somewhat regained before operation
4064	30	2	1	Undelivered -	3	A	—	Mild case
4082	27	1	1	Spontaneous -	FT	A	A	Neurotic
4084	29	6	2a	Spontaneous -	FT	A	A	Well compensated. Mitral stenosis
4090	26	1	2a	Spontaneous (premature)	FT	A	A	Easy labour
4102	47	9	1	Forceps -	8½	A	A	Mild case
4106	24	3	1	Undelivered -	7	A	—	Mild case
4108	27	1	1	Spontaneous -	FT	A	A	Mild case
4132	31	2	1	Undelivered -	7	A	—	Bronchitis, W.R. +
4134	27	3	3	Spontaneous -	7	D	A	Auricular fibrillation. Died
4150	28	1	2a	Spontaneous -	8½	A	A	Mitral incompetence. Well compensated
4237	24	2	2a	Undelivered -	7	A	—	Much improved after 4 weeks rest
4243	18	1	2b	Spontaneous -	8½	A	A	Much improved after ante-natal treatment
4337	20	3	2a	Undelivered -	6	A	—	6 weeks rest. Much improved
4414	30	2	2a	Undelivered -	6	A	—	3 weeks rest. Much improved
4448	21	1	1	Undelivered -	4	A	—	Mild case. Mitral incompetence

B.—Cases sent in as Emergencies.

Reg. No.	Age.	Parity.	Class.	Delivery.	Stage of Pregnancy on Admission.	Result.		Remarks.
						M.	Ch.	
744	22	1	2a	Undelivered -	7	A	—	Mitral stenosis. Improved with rest
763	35	5	2a	Spontaneous -	8½	A	A	Mitral stenosis. Improved with rest
769	38	7	2b	Cæsarean section -	8	D	A	Died on 6th day of puerperium. Severe mitral stenosis
894	39	7	2b	Spontaneous (premature)	FT	A	A	Mitral stenosis. Some loss of compensation
907	21	2	2b	Undelivered -	9	—	—	Auricular fibrillation. Irreg. dismissal
934	25	1	2a	Spontaneous (premature)	6	A	A	2 weeks ante-natal treatment. Readmitted at term
1059	43	9	3	Spontaneous (premature)	7	A	A	Auricular fibrillation. Anasarca. Good recovery
1198	38	5	3	Spontaneous -	8	A	SB	Auricular fibrillation and complete loss of cardiac reserve. Marked improvement on dismissal
1209	26	5	2b	Cæsarean section -	8	A	A	Sterilised. Mitral incompetence. Some failure of compensation
1307	34	7	2a	Cæsarean section -	FT	A	A	Sterilised
1339	26	2	2b	Spontaneous -	9	A	A	Precipitate labour. Good recovery
1429	41	—	2a	Undelivered -	6	A	—	Well compensated. Mitral stenosis
1464	27	3	2a	Spontaneous (premature)	8½	A	A	Marked anæmia
1672	19	1	1	Spontaneous -	FT	A	A	Mild case
1683	29	3	2b	Cæsarean section -	8	A	A	Sterilised. Some loss of cardiac reserve
1691	42	10	2b	Manual extraction	8	A	A	Hypothyroid case. Tachycardia, etc.
1843	30	1	2b	Spontaneous -	FT	A	A	Precipitate labour
2078	27	3	1	Spontaneous -	FT	A	A	Mild case. Anæmia
2345	29	1	1	Forceps -	FT	A	SB	After delivery patient commenced to fibrillate. This ceased after 24 hours and patient made a good recovery
2446	17	1	2a	Spontaneous -	FT	A	A	3 weeks ante-natal treatment. Delivered at term
2516	21	2	2a	Spontaneous -	FT	A	A	Mitral incompetence.
2549	35	4	2b	Spontaneous -	FT	A	A	Mitral stenosis
2563	40	6	2a	Spontaneous (premature)	7½	A	A	Mitral incompetence
2578	20	1	1	Spontaneous (premature)	8	A	A	Mitral incompetence. Mild case
2655	36	7	1	Spontaneous -	FT	A	A	

Reg. No.	Age.	Parity.	Class.	Delivery.	Stage of Pregnancy on Admission.	Result.		Remarks.
						M.	Ch.	
2785	18	2	1	Undelivered -	7	A	—	Anæmia predominant feature
2852	33	4	2b	Cæsarean section -	6	A	A	Sterilised. Double mitral lesion. 8 weeks ante-natal treatment
2872	40	1	1	Undelivered -	6	A	—	Patient had had hæmatemesis outside
3067	28	1	2b	Undelivered -	8	D	—	Sudden death. No p.m. granted
3077	29	4	3	Spontaneous (premature)	7½	A	A	Auricular fibrillation. 4 weeks ante-natal care. Marked improvement on dismissal
3303	24	2	2b	Spontaneous (premature)	8½	A	A	Admitted in labour. Mitral stenosis
3367	27	1	2b	Undelivered -	3	A	—	Mitral stenosis. Right bundle branch block. Irreg. dismis.
3448	25	3	2a	Forceps -	FT	A	A	Mitral stenosis
3950	40	4	1	Undelivered -	7	A	—	Mitral incompetence
3961	36	3	2b	Forceps -	8	A	A	4 weeks ante-natal care. Improved with rest
4129	—	3	2a	Missed abortion -	8	A	—	Mitral incompetence. Some loss of reserve
4212	27	1	1	Spontaneous -	FT	A	A	Mild case
4302	25	2	2a	Spontaneous -	FT	A	A	

C.—Cases not under Ante-natal Supervision.

1071	28	7	1	Forceps -	FT	A	A	Mild case
1220	28	6	2b	Miscarriage -	6½	A	—	Mitral incompetence. Some loss of reserve
2147	28	5	2a	Spontaneous -	7½	A	A	No ante-natal care.
2645	31	2	3	Craniotomy -	FT	A	SB	Auricular fibrillation. Contracted pelvis. Dismissed in much improved state

MATERNAL DEATHS.

There were 119 maternal deaths in the Hospital during the year 1930. Twenty-three of these were in cases which had been under ante-natal supervision, 85 were in cases sent in as emergencies, and the remaining 11 were in cases other than emergencies, which had not been under ante-natal supervision. The causes of death are summarised in the following table :—

Table—Summary of the Causes of Maternal Deaths in 1930.

Cause of Death.	Categ. A.	Categ. B.	Categ. C.	Total Deaths.
Obstetrical complications - - -	3	10	1	14
Puerperal sepsis - - - - -	2	18	2	22
Toxæmia of pregnancy - - - -	4	18	—	22
Ante-partum hæmorrhage - - -	1	12	4	17
Heart disease - - - - -	5	4	—	9
Deaths under anæsthesia - - -	3	1	1	5
Post-partum hæmorrhage - - -	3	10	1	14
Meningitis - - - - -	—	2	—	2
Others (pulmonary embolism, pyelitis, chorea, etc.) - - - - -	2	10	2	14
Totals - - -	23	85	11	119

Since there were 4,469 admissions to the Hospital during the year, the death-rate was 26.6 per thousand (or 2.6 per cent.). The death-rates in the three categories, A, B, and C, were as follows :—

- Category A (cases under ante-natal supervision):
death-rate - - - - - 1.2 per cent.
- Category B (emergency cases sent in as such):
death-rate - - - - - 4.3 per cent.
- Category C (cases not under previous supervision):
death-rate - - - - - 1.6 per cent.

A brief summary of each of the fatal cases is given in the following tables:—

Table—Summary of Fatal Cases during 1930.

Cause of Death.	Reg. No.	Age.	Parity.	Category.	Remarks.
Obstetrical complications	364	39	1	B	Delayed labour. Early rupture of membranes. Repeated examinations before admission. C.P. Elderly primigravida. Os only 3 fingers dilated after 5 days' labour. Sedatives. Three rigors next day. Manual dilatation and craniotomy. P.P.H. Death from shock.
do.	420	40	12	B	Collapsed on admission. Uterus ruptured. No details of labour. Blood transfusion. Hysterectomy. Died next day.
do.	542	24	1	B	C.P. Delayed labour. Cæsarean section. Bronchitis. Cardiac failure. Auricular fibrillation. Died day after delivery.
do.	780	38	7	A	Hydramnios. Delayed labour. Shock. Death after delivery of first child.
do.	1234	27	4	A	Rachitic flat pelvis. Bronchitis. Repeat Cæsarean section. Patient developed post operative anuria and terminal pneumonia. Died on sixth day of puerperium.
do.	2615	36	10	B	Admitted as case of "failed forceps." Os was not fully dilated. Rupture of uterus diagnosed. Death occurred several hours after admission.
do.	2778	48	5	B	Admitted in collapsed condition. Impacted transverse presentation. Outside interference. After delivery, ruptured uterus diagnosed. Hysterectomy performed. Died on day following operation.
do.	3409	23	1	A	Slight degree of contracted pelvis. Low forceps delivery. Patient did not take anaesthesia well and died shortly after operation from what appeared to be obstetric shock.

Cause of Death.	Reg. No.	Age.	Parity.	Category.	Remarks.
Obstetrical complications	3535	28	1	B	Dystocia. Midforceps delivery followed by sudden collapse and death. P.M. examination revealed no abnormality, and death was due to obstetric shock.
do.	3753	34	1	B	Dystocia (48 hours in labour). Generally contracted pelvis. After unsuccessful attempt at forceps delivery, podalic version was performed. The patient collapsed and died a few hours later. P.M. examination showed rupture of uterus.
do.	3797	26	4	C	Rachitic pelvis, slight contraction. After attempt at forceps delivery, craniotomy was performed. The uterus was found to be ruptured and the patient died four days later. P.M. examination showed generalised peritonitis.
do.	3968	48	2	B	Dystocia marked, craniotomy performed. Pneumonia developed and patient died on 12th day of puerperium. P.M. examination confirmed clinical findings.
do.	4252	38	6	B	Dystocia on account of hydrocephalus. Admitted with (?) ruptured uterus. After delivery, laparotomy was performed and the ruptured uterus repaired. Patient died in spite of blood transfusion.
do.	4295	43	4	B	Admitted as case of "failed forceps" outside. Markedly contracted rachitic pelvis. Cæsarean section performed. Patient died 4 days afterwards with acute pulmonary œdema. No. P.M. examination granted.
Puerperal sepsis	170	54	1	B	B.B.A. Retained placenta. P.P.H. Febrile. Antistreptococcal serum. Puerperal fever. Died 11th day after admission.
do.	21	26	3	B	Breech. Hydrocephalus and meningocele. Perforation. Difficult delivery. Febrile from 3rd day. Death from sepsis on 10th day.
do.	238	36	9	B	Incomplete abortion. Too ill for operation at first. Febrile from 5th day, uterus cleared out. Pulmonary embolism next day. Died 7th day after admission. P.M., pelvic cellulitis and thrombosis. Hæmatoma (? traumatic) right broad ligament. Thrombosis and infarction of lung.

Cause of Death.	Reg. No.	Age.	Parity.	Category.	Remarks.
Puerperal sepsis	3541/29	20	1	B	Brecch with extended legs. Attempted delivery by traction before admission. Extensive vaginal lacerations. Manual delivery. Antistreptococcal serum. Febrile from 2nd day. Septicæmia. Died 35th day of puerperium.
do.	177	26	1	A	Hyperemesis till 7th month. Later œdema, visual disturbance, headache, albuminuria. B.P. 160/118. Induced (bougies). Febrile from 2nd day. P.P.H. Antistreptococcal serum. Died 4th day of puerperium.
do.	291	25	1	B	C.P. Forceps delivery. Albuminuria and toxæmia. Pyuria. Febrile puerperium. Sudden death on 15th day from acute cardiac failure with pulmonary œdema—probably embolism.
do.	302	37	6	B	A.P.H. Placenta prævia. Intermittent bleeding for 6 weeks. Spontaneous delivery after severe hæmorrhage. Manual removal of placenta. Febrile—septicæmia. Died 2 months later. P.M., extensive abscesses in pelvis and in psoas muscles.
do.	929	41	8	B	Pre-eclamptic toxæmia. Induction (bougies). Toxæmia cleared up after delivery but sepsis developed. Died 13th day. P.M. Primary peritonitis. No uterine sepsis seen.
do.	1167	24	4	C	Admitted to ante-natal wards with hyperemesis gravidarum and pyelitis. Dismissed well after 5 weeks. Readmitted in labour in rather poor condition. Died on 19th day of puerperium from puerperal septicæmia.
do.	1240	39	6	A	Admitted on account of prolapsed hypertrophied cervix which was ulcerated and septic. Five days later was delivered of hydrocephalic macerated foetus (no instrumental interference). Serum given. Died on 11th day of puerperium from septicæmia.
do.	1248	36	7	B	Delivered on day of admission. Placenta adherent and manually removed. Serum given. On 4th day of puerperium developed phlegmasia followed by pyæmia from which she died 9 weeks later.

Cause of Death.	Reg. No.	Age.	Parity.	Category.	Remarks.
Puerperal sepsis	1351	26	4	B	Placenta prævia (central). Outside examination by midwife and doctor. Version performed in hospital. Died eleven days after admission from puerperal septicæmia.
do.	1360	31	3	B	Twin pregnancy. Dystocia. Outside interference (failed forceps). Forceps delivery in hospital. Septicæmia and peritonitis. Died 4 days after admission.
do.	1938	27	3	C	Admitted exsanguinated with central placenta prævia. Abdominal hysterotomy operation performed. Developed broncho-pneumonia ten days after operation and died on the 13th day. P.M. showed uterine sepsis.
do.	2389	37	8	B	Admitted with incomplete abortion. Debilitated and exsanguinated. Developed puerperal sepsis 5 days after digital curettage. Died 10 days later.
do.	2514	21	1	B	Dystocia but spontaneous delivery. Retained placenta — manually removed. Never febrile. Died on 5th day of puerperium. P.M., Puerperal sapræmia, pelvic cellulitis and bilateral hydronephrosis.
do.	2789	33	6	B	Incomplete abortion. Digital curettage. Death 10 days later from puerperal septicæmia.
do.	3292	25	1	B	Cæsarean section for rachitic pelvis after prolonged labour. Peritonitis developed along with hypostatic pneumonia. Death on 5th day of puerperium.
do.	3101	25	1	B	Cæsarean section for rachitic pelvis after delayed labour. Sepsis appeared in wound, which repeatedly broke down. Faecal fistula developed. Patient died after 30 days in hospital.
do.	3805	26	4	B	Cæsarean section after "failed forceps outside." Marked rachitic deformities. The wound became septic. Patient died of generalised peritonitis 3 days later.
do.	3592	29	1	B	Admitted as case of "failed forceps" outside. Craniotomy performed in hospital. Anti-streptococcal serum and blood transfusion. Died 12 days after admission.

Cause of Death.	Reg. No.	Age.	Parity.	Category.	Remarks.
Puerperal sepsis	4386	19	1	B	Admitted as case of "failed forceps" outside. Lacerated on admission. Delivered two days later by forceps. Died on 3rd day of puerperium from puerperal septicæmia.
Toxæmia of pregnancy	35	21	1	B	Three eclamptic fits before admission in coma. Twelve further fits. B.P. 230/200. Gastric lavage. Veratrone. Delivery with low forceps. Acute pulmonary œdema on 3rd day of puerperium.
do.	158	25	1	B	Vomiting for 6 weeks. Exhaustion and toxæmia. Cerebral symptoms before death 10 days after admission.
do.	1023	30	1	A	Pyelitis early in pregnancy. Improved. Fulminating ante-partum eclampsia. 30 fits. Died 13 hours after admission.
do.	1039	35	3	B	Admitted in debilitated state suffering from hyperemesis gravidarum. Patient aborted completely but failed to improve thereafter. Anæmia (secondary) and toxic symptoms were severe. She died after having been in hospital for three months.
do.	1455	40	1	B	Pre-eclamptic toxæmia. Surgical induction (twice). Patient miscarried but became more toxic. Jaundice appeared and patient died on 7th day of puerperium.
do.	1466	33	1	B	Admitted after delivery at home, suffering from post-partum eclampsia. Patient took eleven seizures and died on the day after admission.
do.	1572	36	1	B	Admitted comatose, several fits before admission. Severe fits after admission. Cæsarean section performed. Died from cerebral hæmorrhage the day after admission.
do.	2205	26	1	B	Patient admitted in coma (severe eclampsia). Fits continued despite treatment. Patient died on day of admission undelivered.
do.	2467	32	2	A	Admitted with eclampsia, delivered shortly thereafter. Developed uræmia and died three days later.
do.	2511	18	1	B	Admitted with intra-partum eclampsia, severe type, delivered on the same day. Died on the following day.

Cause of Death.	Reg. No.	Age.	Parity.	Category.	Remarks.
Toxæmia of pregnancy	2664	36	3	B	Admitted moribund, severe eclampsia, died shortly after admission undelivered. Post-mortem confirmed the diagnosis.
do.	2820	28	2	B	Admitted on account of chronic nephritis. Miscarriage followed by severe abdominal pain and shock from which she died. Post-mortem revealed large retro-peritoneal hæmorrhage surrounding the left kidney of unknown origin.
do.	2932	40	7	B	Admitted comatose with antepartum eclampsia. Delivered spontaneously. Coma persisted and death occurred several hours after admission.
do.	3151	24	1	B	Admitted with severe hyperemesis gravidarum. Abdominal hysterotomy on day after admission. Died on day after operation.
do.	3224	28	3	B	Suffered from pre-eclamptic toxæmia. Admitted after delivery of twins outside. Died several hours after admission. P.M. examination confirmed diagnosis.
do.	3392	40	8	A	Was in antenatal wards with severe albuminuria. Left hospital irregularly. Readmitted moribund. Died during induction of anæsthesia for a forceps delivery four days after readmission.
do.	3415	27	3	A	Admitted to antenatal wards with severe pyelitis. Induction of abortion performed. She died on 9th day of puerperium. P.M. examination showed pyelitis and multiple abscesses in kidneys.
do.	3664	34	3	B	Eclampsia (1 fit prior to admission). After a spontaneous and complete miscarriage, patient developed pulmonary œdema and died 4 days after admission.
do.	3736	23	2	B	Admitted as eclampsia. Birth occurred outside. Comatose. Suppression of urine. Died five days after admission.
do.	3771	45	1	B	Severe pre-eclamptic toxæmia at home. Admitted with fulminating eclampsia. Died 36 hours after admission.

Cause of Death.	Reg. No.	Age.	Parity.	Category.	Remarks.
Toxæmia of pregnancy	3821	31	2	B	Treated successfully in antenatal wards for hyperemesis gravidarum. Dismissed irregularly. Readmitted seriously ill. Abdominal hysterotomy immediately performed, but death occurred shortly after operation.
do.	3826	26	1	B	Eclampsia. 3 fits before admission. 10 fits after admission. Died 7 hours after admission. P.M. examination showed cartical neerosis of kidney.
Ante-partum hæmorrhage	310	40	13	B	Accidental hæmorrhage, concealed and apparent. Repeated bleeding. Spontaneous delivery. Death from shock.
do.	535	37	10	B	A.P.H. Moribund on admission. Died a few hours later. No P.M.
do.	406	39	4	A	Accidental hæmorrhage, concealed and apparent. Died a few hours after admission.
do.	667	41	8	B	Placenta prævia. Died 3 hours later. Moribund on admission.
do.	665	40	7	B	Incomplete abortion. Severe bleeding. Died a few hours later.
do.	538	—	10	B	A.P.H. Died 30 minutes after admission. No P.M.
do.	987	28	5	C	Hyperemesis. Induction proposed but lateral placenta prævia diagnosed. A.P.H. Packing. Spontaneous delivery. Died 1 hour later.
do.	2456	35	9	B	Admitted from "district." Severe accidental hæmorrhage (concealed). Profound shock. Died soon after admission. Confirmed diagnosis at P.M.
do.	2488	37	3	B	Admitted as pre-eclamptic toxæmia. Went into labour five days later. Severe accidental hæmorrhage (mixed) profound shock. Died undelivered.
do.	2617	33	7	B	Admitted on account of accidental hæmorrhage. Spontaneous delivery. 3 days later patient suddenly developed puerperal mania and jumped from a window, committing suicide.
do.	2646	42	10	B	Admitted with concealed accidental hæmorrhage. Delivered after medical induction. Post-partum hæmorrhage. Blood transfusion. Died 2 days later.
do.	2672	31	6	C	Admitted exsanguinated. Central placenta prævia. Delivered after version. Died on day of admission.

Cause of Death.	Reg. No.	Age.	Parity.	Category.	Remarks.
Ante-partum hæmorrhage	2797	32	5	C	Admitted on account of mixed accidental hæmorrhage. Forceps delivery and gum acacia transfusion. Patient died 3 hours after admission.
do.	2815	39	7	B	Admitted with toxic symptoms. Septic eruptions on body generally. Accidental hæmorrhage on day following admission from which she died. Post-mortem confirmed diagnosis and showed chronic mitral stenosis and acute endocarditis.
do.	2868	30	6	B	Admitted from the "district" with placenta prævia. Patient died from a combination of hæmorrhage and obstetric shock a few hours after admission.
do.	3241	40	5	B	Central placenta prævia. Classical Cæsarean section performed. On 6th day patient suddenly collapsed and died. No cause could be ascertained. No P.M. granted.
do.	3667	35	8	C	Admitted with accidental hæmorrhage (mixed). After spontaneous delivery became deeply jaundiced. Severe hæmatemesis. Died 2 days after admission. P.M. examination showed toxic jaundice and ante-mortem digestion of œsophagus.
Deaths under anæsthesia	1855	32	1	A	Dystocia and occipito posterior position. Patient collapsed after delivery by low forceps. Death due to chloroform poisoning.
do.	1933	20	3	C	While under anæsthesia for the digital curettage of an incomplete abortion the patient suddenly collapsed and died. Death was due to chloroform poisoning.
do.	3280	25	3	A	Cæsarean section performed for rachitic flat pelvis. Patient died under the anæsthetic.
do.	3706	?	1	B	This patient died of delayed chloroform poisoning. She had a prolonged chloroform anæsthetic outside and was admitted as "failed forceps." In hospital, forceps delivery was successful under spinal anæsthetic. Died on 4th day of puerperium. P.M. examination showed signs of delayed chloroform poisoning.
do.	4171	33	4	A	Case of missed abortion. Died under anæsthesia (ether and oxygen) while uterus was being emptied.

Cause of Death.	Reg. No.	Age.	Parity.	Category.	Remarks.
Post-partum hæmorrhage	303	33	5	B	B.B.A. Retained placenta. P.P.H. Died 4 hours after admission.
do.	340	27	1	B	B.B.A. Retained placenta. P.P.H. Collapsed. Died 3 hours after admission.
do.	512	41	7	B	Accidental hæmorrhage at home. Admitted on account of severe post-partum hæmorrhage. Died 2 hours after admission.
do.	570	34	2	A	Hydramnios. Spontaneous delivery. P.P.H. Retained placenta. Further P.P.H.
do.	599	27	1	B	B.B.A. Retained placenta. P.P.H. Manual removal. Died 3 hours later.
do.	743	21	1	B	B.B.A. (twins). Retained placenta. P.P.H. Died on admission.
do.	2802	29	1	B	Admitted on account of post-partum hæmorrhage, with retained placenta. Baby born outside. Placenta manually removed. Died 3 hours after admission.
do.	2830	32	6	A	Admitted as inevitable abortion in poor condition. The patient aborted, but bleeding continued even after digital curettage. Patient died 2 days after admission
do.	2914	32	9	B	Admitted collapsed after complete abortion. Despite treatment and blood transfusion, patient died on following day.
do.	2961	36	?	B	Admitted on account of retained placenta and post-partum hæmorrhage. Placenta manually removed. Death occurred 1½ hours after admission.
do.	3053	33	9	C	Twin pregnancy. Severe hæmorrhage after birth of first twin. This continued after birth of second twin. Placenta manually removed. Blood transfusion. Death several hours after delivery.
do.	3758	22	3	B	Admitted on account of retained placenta and post-partum hæmorrhage. She died 1 hour after admission. P.M. examination showed densely adherent placenta and pyelitis.
do.	3806	30	---	A	Admitted to ante-natal wards in poor condition, having had severe hæmatemesis. Slight post-partum hæmorrhage. She died shortly after delivery. P.M. examination showed chronic duodenal ulcer and some healed pulmonary tuberculosis.

Cause of Death.	Reg. No.	Age.	Parity.	Category.	Remarks.
Post-partum hæmorrhage	4098	33	8	B	Admitted as incomplete abortion, exsanguinated and eollapsed. Digital curettage was performed. Patient died 3 hours after admission.
Heart disease	769	—	7	B	Cardiac failure. Cæsarean section (spinal anæsthesia). Improved for a time. Wound gave way during coughing. Sudden death from acute cardiac dilation and pulmonary œdema on 7th day of puerperium.
do.	745	38	13	A	Chronic renal disease and myocarditis. Four weeks in ante-natal wards. Complete miscarriage. Death on 4th day of puerperium.
do.	764	38	4	A	Admitted with advanced cardiac disease (mitral stenosis). Cæsarean section performed. Died three weeks later from cardiac failure.
do.	2733	27	4	A	Mitral stenosis and loss of compensation. Delivered spontaneously 15 days after admission. During puerperium cardiac decompensation became worse. Died on 8th day of puerperium.
do.	3009	35	6	A	Admitted with advanced mitral stenosis. Treated in ante-natal wards for four weeks. Cæsarean section at term. Died of cardiac failure 6 hours after operation. P.M. examination confirmed diagnosis.
do.	3067	28	1	B	Admitted to ante-natal wards with severe mitral stenosis. Compensation was lost and became succeedingly worse despite treatment. Patient died undelivered 12 days after admission.
do.	3073	37	5	A	In ante-natal with albuminuria of pregnancy. Readmitted with pre-eclamptic toxæmia. Developed subacute endocarditis after spontaneous delivery. Died 1 month later.
do.	3618/29	19	1	B	Auricular fibrillation and cardiac failure on admission. Cæsarean section 10 days later. Fairly well for a time but progressive cardiac decompensation. Died 5 weeks after delivery.
do.	3783	29	5	B	Treated in ante-natal wards for cardiac disease and bronchitis. Readmitted shortly after birth of child. Developed severe bronchitis and basal pneumonia. Died 8 days after admission.

Cause of Death.	Reg. No.	Age.	Parity.	Category.	Remarks.
Malignant cachexia	109	40	6	B	Carcinoma of breast. Malignant cachexia. Induction. Died 6th day of puerperium. P.M. Carcinoma of breast and generalised carcinomatosis.
Pyelitis	3791/29	21	1	B	Pyelitis. Improved but relapsed. Toxæmia and jaundice. Surgical induction. Died from toxæmia on 8th day of puerperium.
Chorea	220	23	2	A	Chorea (recurrent). Cardiac symptoms. Induction. Died undelivered. P.M. Acute endocarditis.
Cerebral tumour	150	27	2	B	Hyperemesis. Jacksonian fits. Spontaneous delivery. Died 16 days later. P.M. Glioma of right cerebral hemisphere.
Tumour of bowel	1031	44	9	B	Transverse presentation. Delayed labour. Obstetrical shock. Twin pregnancy. Died after delivery of first child. P.M. Carcinoma of sigmoid with perforation and general peritonitis. Extensive carcinomatosis of liver.
Anæmia	767	39	5	B	Pyelitis. Albuminuria. Premature delivery (twins). Epistaxis. Progressive anæmia. Died 1 month later.
Carcinoma of cervix	3385/29	35	10	A	Inoperable carcinoma of cervix. Cæsarean section at term. Died 3 days later (peritonitis).
Malnutrition	1002	35	7	C	B.B.A. Sent in collapsed. Asthenia and malnutrition. Purpuric spots. Anæmia. Died 8th day.
Meningitis	1686	33	1	B	Patient admitted in deep coma as eclamptic. Died five hours later. Post-mortem revealed empyema of gall bladder and generalised purulent meningitis. Undelivered.
do.	4350	42	9	B	Admitted on account of otitis media, which developed into a basal meningitis, from which patient died 7 days after admission. P.M. examination confirmed diagnosis.
Broncho-pneumonia	—	30	—	B	Admitted with broncho-pneumonia. Delivered by low forceps on the same day. The pneumonia became generalised and she died on the 4th day of the puerperium.
Ulcerative colitis	3488	29	5	B	Admitted as antenatal case with marked ulcerative colitis. No bacteriological evidence of typhoid paratyphoid or dysentery. Became progressively worse after delivery and died 5 days after admission.

Cause of Death.	Reg. No.	Age	Parity.	Category.	Remarks.
Antepartum pyæmia	3149	24	1	B	Admitted on account of dystocia with history of dyspnœa and vomiting of several weeks' duration. Died shortly after forceps delivery. P.M. examination showed pulmonary embolism, pulmonary abscesses, septic infarction of heart and pyelonephritis.
Gangrene of lung	2262	22	3	B	Admitted on account of post-partum hæmorrhage. Placenta manually removed. Developed pneumonia on 10th day of puerperium. Empyema followed, and finally patient died from what was found at post-mortem to be gangrene of lung.
Pulmonary embolism	2931	30	3	C	Contracted pelvis. Classical Cæsarean section performed. Progress satisfactory until 5th day of puerperium when death occurred suddenly with signs of pulmonary embolism.
Ectopic pregnancy	2953	22	1	B	Ectopic pregnancy (advanced), with infection of the sac was found at operation. Patient died of peritonitis on the 17th day after operation.

STILLBIRTHS AND NEONATAL DEATHS.

1. *Stillbirths.*

The total number of stillbirths in the indoor cases during the year was 396. Of these, 83 were in cases in Category A, 270 in cases in Category B, and 43 in cases in Category C. Since the total number of viable children born was 2,837, this gives an incidence of stillbirth of 13.9 per cent.

In the Outdoor Department the total number of stillbirths was 85, giving an incidence rate of 2.1 per cent.

2. *Neonatal Deaths.*

The total number of neonatal deaths was 160. Of these, 46 were in cases in Category A, 79 in cases in Category B, and 35 in cases in Category C. The number of viable children born being 2,837, this gives an incidence of neonatal death of 5.6 per cent. In the three Categories A, B, and C, the incidence rates were 3.7, 7.4, and 6.3 respectively; 84 of the neonatal deaths occurred during the first day (=52.5 per cent. of all neonatal deaths). If these cases are deducted from the total, there remain 76 cases, giving a neonatal death-rate of 2.6 per cent.

PUERPERAL PYREXIA.

The following table presents the figures for the year ending 1930, and includes cases notified after dismissal from hospital which were regarded as being primarily hospital cases:—

Annual Return Figures for Puerperal Fever and Pyrexia for year ending 1930.

		Total Cases.		Cases per 1,000 Births.		Deaths.		Deaths per 1,000 Births.		Case Mortality per cent.	
		Fever.	Pyrexia	Fever.	Pyrexia	Fever.	Pyrexia	Fever.	Pyrexia	Fever.	Pyrexia
Indoor	-	130	63	45.3	21.9	24	7	8.4	2.4	18.5	11.1
Outdoor	-	49	24	12.3	6.0	6	1	1.5	.3	12.2	4.2

The above table was culled from the Public Health statistical records and checked from the Hospital temperature charts. It will be seen that no attempt has been made to classify the patients in their categories.

Without attempting to differentiate between Puerperal Sepsis and Puerperal Pyrexia it was possible to ascertain from the records of the total number cases notified as one or other condition. That 30.1 per cent. were in Category A, 56.4 per cent. in Category B, and 8.3 per cent. in Category C. Attention is directed to the relatively high percentage in Category B (the cases sent in as emergency cases).

ABORTIONS.

There were 822 cases admitted on account of incomplete, threatened, or missed abortion during the year. Of these, 151 were in Group A, 592 in Group B, and 179 in Group C: 120 of the patients were primigravidæ and the remaining 702 were multiparæ.

MOLE.

There were 15 cases of mole in the Hospital.

APPENDIX.

1.—STOBHILL CASES.

During the year, 778 cases were transferred on admission to the beds reserved for the Maternity Hospital at Stobhill and other Corporation Hospitals. Of this total, 750 were delivered. The number of children born was 750, of which 727 were livebirths and 23 were stillbirths. There were 6 maternal deaths. The incidence of stillbirth was 3·0 per cent. The number of cases recorded as normal was 732, and the number recorded as abnormal was 46 (=5·9 per cent.).

2.—DISTRICT CASES.

During the year, 4,079 cases were attended in their own homes in the district. The number delivered was 3,812, and 267 aborted or were undelivered. The number of live-born children was 3,812, and there were 85 stillbirths. There were 5 maternal deaths, and the incidence of stillbirth was 2·1 per cent. The number of cases recorded as normal was 3,713, and the number recorded as abnormal was 366 (8·9 per cent.).

The number of visits paid to patients in the Outdoor (District) Department was 44,894 (an average of 3,741 visits per month).

